



CHARACTERIZATION, DIGESTIBILITY AND PREBIOTIC EFFECT OF THE FRUIT EXTRACT OF *Stenocereus queretaroensis* F.A.C. WEBER BUXBAUM

CARACTERIZACIÓN, DIGESTIBILIDAD Y EFECTO PREBIOTICO DEL EXTRACTO DEL FRUTO DE *Stenocereus queretaroensis* F.A.C. WEBER BUXBAUM

José J. Gaytán-Andrade, Luis E. Cobos-Puc, Anna Iliná,
Illeana M. M. Moreno-Dávila, Cristóbal N. Aguilar and Sonia Y. Silva-Belmares*

Autonomous University of Coahuila, School of Chemistry, Postgraduate Program in Food Science and Technology, Research Group of Chemist Pharmacologist, Saltillo, Coahuila, Mexico.

*Corresponding author (yesenia_silva@uadec.edu.mx)

SUMMARY

Digestibility is the ease with which food is converted in the digestive system into useful substances for nutrition; it comprises the hydrolysis of complex molecules and the absorption of small molecules in the intestine. The microbiota is essential in these processes and the malfunction of these microorganisms is associated with metabolic diseases. The objective of this study was to characterize and evaluate the digestibility and prebiotic effect of the *Stenocereus queretaroensis* fruit extract, since its nutritional value and functionality are unknown. The extract was obtained from the fruit at 4 °C using a juice extractor; then, it was filtered and lyophilized, and its partial nutritional value and quality were determined by physicochemical and proximal analyses. In addition, techniques were established to evaluate the oral, gastric, and intestinal digestibility, and to evaluate the prebiotic effect. The extract displayed a red color ($L^* = 17.21 \pm 0.1$, $a^* = 38.61 \pm 0.1$, $b^* = 10.60 \pm 0.2$), with a pH of 3.78 ± 0.1 , without crude fat, and 0.77 ± 0.0 % (w/w) of raw fiber, 1.36 ± 0.0 % protein, 88.82 ± 0.8 % of nitrogen-free extract, and prebiotic effect to *Bacillus coagulans*, *Lactobacillus sporogenes* and *L. delbrueckii*.

Index words: *Stenocereus queretaroensis*, characterization, digestibility, prebiotic evaluation.

RESUMEN

La digestibilidad es la facilidad con la que los alimentos se convierten en el sistema digestivo en sustancias útiles para la nutrición; ésta comprende la hidrólisis de moléculas complejas y la absorción de moléculas pequeñas en el intestino. La microbiota es fundamental en estos procesos y el mal funcionamiento de estos microorganismos está asociado con enfermedades metabólicas. El objetivo de este estudio fue caracterizar y evaluar la digestibilidad y el efecto prebiótico del extracto de fruto de *Stenocereus queretaroensis*, ya que se desconocen su valor nutricional y funcionalidad. El extracto se obtuvo del fruto a 4 °C usando un extractor de jugo; posteriormente, se filtró y liofilizó; además, se determinó su valor y calidad nutricional parcial mediante análisis fisicoquímicos y proximales. Asimismo, se establecieron técnicas para evaluar la digestibilidad oral, gástrica e intestinal y para evaluar el efecto prebiótico. El extracto mostró un color rojo ($L^* = 17.21 \pm 0.1$, $a^* = 38.61 \pm 0.1$, $b^* = 10.60 \pm 0.2$), con un pH de 3.78 ± 0.1 , sin grasa cruda y 0.77 ± 0.0 % (p/p) de fibra cruda, 1.36 ± 0.0 % de proteína, 88.82 ± 0.8 % de extracto libre de nitrógeno y efecto prebiótico para *Bacillus coagulans*, *Lactobacillus sporogenes* y *L. delbrueckii*.

Palabras clave: *Stenocereus queretaroensis*, caracterización, digestibilidad, evaluación prebiótica.

INTRODUCTION

Digestibility is a way to determine the utility of foods; that is, the ease with which these are converted in the digestive system into useful substances for nutrition (Yoon *et al.*, 2019). It comprises two processes, the digestion that corresponds to the hydrolysis of complex food molecules and the absorption of small molecules (amino acids, fatty acids) in the intestine (Ye *et al.*, 2018). Microbiota is essential in these processes since the malfunction of these microorganisms is associated with metabolic diseases (Avolio *et al.*, 2020).

A current tendency of the world population is focused on consuming natural foods derived from plants with high nutritional value (Román-Cortés *et al.*, 2018; Orozco-Sifuentes *et al.*, 2019; Ramírez-Jaspeado *et al.*, 2020) and prebiotic properties (Talebi *et al.*, 2020). *Stenocereus queretaroensis* (Cactaceae) is a plant endemic to Mexico (Campos-Rojas *et al.*, 2011). There are different types, but the red one has a higher production and impact on the Mexican economy (Quiroz-González *et al.*, 2018). The Mexican population consumes the pulp (Gaytán-Andrade *et al.*, 2019; Quiroz-González *et al.*, 2018) and the pigmented juice is used to prepare traditional ice drinks (Soto-Castro *et al.*, 2019). Mexico produced 5807 t in the year 2019 (SIAP, 2020).

Stenocereus queretaroensis presents total soluble solids content of 16 °Brix and total sugars of 10 to 11 %. The titratable acidity percentage (citric acid) ranges from 0.15 to 0.50 % and the pH from 3.9 to 5.2, depending on the pitayo type (Gaytán-Andrade *et al.*, 2019).

The *S. griseus* fruit has a pH higher than 4.5 and contains 9.9 % of total sugars (Chuck-Hernández *et al.*, 2016). To date, no extraction methods have been described for the fruit, but ultrasound extractions have been used to extract

active ingredients from Cactaceae (Ramli *et al.*, 2014); however, these methods can modify the integrity of the chemical structure of the components (Sasidharan *et al.*, 2011). On the other hand, the methods for evaluating *in vitro* digestibility of food depend on the type of sample (Bornhorst and Singh, 2014; Brodkorb *et al.*, 2019). Digestibility data of plant extracts are scarce, including those for *S. queretaroensis*; however, simulated *in vitro* oral (using saliva), gastric, and intestinal (small intestine and large intestine) digestibility tests are commonly used for this purpose (Chen *et al.*, 2018).

To date the nutritional and functional information of *S. queretaroensis* extracts is not completely known (Soto-Castro *et al.*, 2019), even so, it could represent a source for development of new functional foods (Gaytán- Andrade *et al.*, 2019). For this reason, the objective of this study was to characterize and to evaluate digestibility and prebiotic property of the extract of red *S. queretaroensis* fruit.

MATERIALS AND METHODS

Plant material and extraction

Red fruits of *S. queretaroensis* (FAC Weber) Buxbaum were harvested in Amacueca, Jalisco, México (coordinates: 19° 56' N, 103° 04' W, altitude of 1361 masl) between May and June 2017 at physiological maturity suitable for consumption. A sample of the plant was identified and deposited at the herbarium of the Antonio Narro Autonomous Agrarian University with the registration number 99186. The extract of the fruit (2.5 kg) without pericarp was carried out in an extractor (Turmix, Mexico City) inside a cold room (4 °C) to guarantee the integrity of the compounds. The extract was centrifuged at $2200 \times g$ for 30 min, filtered with Whatman # 4 paper, and clarified with Whatman G/FA; then, it was dried with a lyophilizer (Labconco, Freeze dryer 5, Kansas, EUA). The yield percentage Y (%) was calculated by gravimetric analysis from the fruit weight (g) used for the extraction as described in the formula:

$$Y (\%) = \frac{\text{weight of the recovered extract (100)}}{\text{weight of fruit}}$$

The extract was stored at -20 °C until use (Hernández *et al.*, 2017).

Physicochemical and proximal analyses

The total soluble solids (TSS, °Brix) were measured using a refractometer (Atago®, PAL-1, Tokyo), pH was determined with a potentiometer (Hanna, pH 209, Plopiș, Romania) and color (L*, a* and b*, using the

CIELAB scale) was measured by using a colorimeter (PCE instrument, CSM-2, Albacete, Spain). These parameters were evaluated in a 1 mg mL⁻¹ solution prepared from the lyophilized extract. °Brix were measured with 100 μL, while pH and color with 50 mL. Additionally, titratable acidity (TA) was measured as the % of citric acid using method 942.15 (AOAC, 2000; Sadler and Murphy, 2010). The test was carried out by titrating 5 mL of a dilution of the lyophilized extract to 0.2 mg mL⁻¹ with 0.1 N NaOH, using 1 % phenolphthalein in ethanol as indicator. The proximal analysis was performed using the AOAC official methods (AOAC, 2000). The moisture and ash content were measured by methods 925.09 and 923.03 using 1 g of lyophilized extract. Ash content was determined by incineration in a muffle at 550 °C. The fat, crude fiber and protein contents were determined by methods 920.39 (Soxhlet), 962.09 and 979.09 (Kjeldahl, N × 6.25) in 10 g of lyophilized extract. Protein content was measured from the previously defatted lyophilized extract. The percentage of the nitrogen-free extract calculated as % Nitrogen-free extract = 100 - (% humidity + % ash + % fat + % crude protein + % crude fiber). The parameter corresponds to the total content of digestible carbohydrates, vitamins and soluble non-nitrogenous organic compounds (Dick *et al.*, 2019). All measurements were on dry weight (DW) basis, except the moisture.

Digestibility

Oral digestibility was measured using the method described by Chen *et al.* (2018) with some modifications. Three healthy 30 to 40-year-old volunteers, without receiving any antibiotic treatment for at least three months, supplied saliva and it was used immediately without any treatment. The test was done in two tubes, in the first one 1 g of extract was placed and in the other 1 g of distilled water was used as a control; then, 2 mL of saliva were added to each tube, mixed for 10 s and incubated at 37 °C for 2 min. Right away, the amylase was inactivated in a boiling water bath for 5 min. The tubes were centrifuged at $5000 \times g$ for 10 min to remove the supernatant and the precipitate was dried at 50 °C for 3 h. Finally, the precipitate was weighed and the percentage of oral digestion calculated using the formula:

$$\% \text{ Oral digestibility} = 100 - \left[\frac{(\text{g precipitate})}{\text{g extract}} \right]$$

Gastric digestibility percentage was measure using a combination of the methods described by Chen *et al.*, 2018) and Mkandawire *et al.* (2015) with some modifications. Solution A was prepared by mixing an electrolyte solution (3.1 g of NaCl, 1.1 g of KCl, 0.25 g of CaCl₂ and 0.6 g of NaHCO₃ in 1 L of distilled water) with a solution of 0.1 mol

HCl L⁻¹ at 1:1 ratio. Solution B was prepared with 1.5 mg mL⁻¹ pepsin dissolved into 0.1 mol L⁻¹ sodium phosphate buffer, pH 3. All solutions were dissolved at 37 °C. Gastric digestibility was tested in a tube containing 1 g of the lyophilized extract along with 1 g of distilled water as a control. First, gastric fluid was prepared with 7 mL of solution A and 3 mL of solution B, adjusted to pH 3 with 0.1 mol L⁻¹ HCl; then, 10 mL of gastric fluid were added to each tube to start digestion and stirred for 2 h at 37 °C. Pepsin was inactivated in a boiling water bath for 5 min, the tubes were centrifuged at 5000 × *g* for 10 min, and the supernatant removed.

The precipitate was washed with 20 mL of 0.1 mol L⁻¹ sodium phosphate buffer (pH 7) and recovered using centrifugation. The precipitate was dried for 3 h at 50 °C and weighed. The percentage of gastric digestion was calculated using the formula

$$\% \text{ Gastric digestibility} = 100 - [(g \text{ precipitate}) \times 100 / g \text{ extract}]$$

Digestion in the large intestine was investigated by *in vitro* fermentation with the Daisy[®] Method as described by Coblenz *et al.* (2019) with some modifications. Before starting the test, solutions A, B, and fecal inoculum were prepared. Solution A was prepared with 10 g of KH₂PO₄, 0.5 g of MgSO₄ · 7H₂O, 0.5 g of NaCl, 0.5 g of urea, and 0.1 g of CaCl₂ · 2H₂O in 1 L of distilled water. Solution B was prepared with 15 g of Na₂CO₃ and 1 g of Na₂S · 9H₂O in 1 L of distilled water. The human fecal inoculum solution was prepared from a mixture of equal amounts of stools donated by two healthy volunteers without antibiotic treatment. The solution was prepared with 200 g of feces dissolved in 2 L of sterile 0.9 % saline solvent at 37 °C. The solution was homogenized using a glass rod and filtered through a 0.15 mm pore mesh. The tests were carried out in 4 × 4 cm polyester-polyethylene bags (filter bag F57) with a pore size of 25 µm. The bags were rinsed using acetone and air-dried. The empty bag was weighed, and then 1 g of extract was placed inside. In addition, an empty bag was weighed and used as a control and all bags were heat-sealed. All the glassware and the digestion jar were kept at 37 °C.

Simultaneously, a digestion solution was prepared with 400 mL of human fecal inoculum solution, 1.2 L of solution A and 250 mL of solution B, and maintained at 37 °C. The samples were evenly distributed in the jar in which the digestion solution was added; then, the jar was covered, purged, supplied with CO₂ for 30 s and incubated for 24 h at 37 °C in the Daisy[®] incubator. At the end of the incubation, the jar was removed and the liquid drained. The digested bags were recovered, rinsed with distilled water until the

water was clean; then, the microbial residues and any remaining soluble fractions were removed with a neutral detergent solution. Subsequently, they were dried for 3 h at 50 °C and weighed. Calculation was performed as follows:

$$\text{Intestinal digestibility} = 100 - [(g \text{ bag of digested extract} - g \text{ bag control}) \times 100 / g \text{ of undigested extract}]$$

Prebiotic evaluation

Commercial strains of *B. coagulans*, *L. sporogenes*, *B. clausii*, *L. fermentum* and *L. delbrueckii* were used in the assay. The strains were activated using MRS 55 g L⁻¹ broth at 37 °C for 12 h. The prebiotic effect was evaluated through bacterial growth, by combining the McFarland technique and the growth kinetics of each strain during the exponential phase (De *et al.*, 2014; Jurado-Gámez *et al.*, 2014). The growth of each strain at 24 h was compared to evaluate the prebiotic effect. The absorbance of the McFarland 0.5 scale was used to adjust the bacterial suspension of the initial inoculum (1 × 10⁸ CFU mL⁻¹) used in each growth kinetics. The growth kinetics of each strain was performed in tubes with 9 mL of MRS, 1 mL of *S. queretaroensis* extract at 1000 µg mL⁻¹ and 300 µL of bacterial suspension as initial inoculum. A blank without inoculum was used to adjust the absorbance. Tubes were incubated at 37 °C for 24 h in anaerobiosis, and absorbance was measured every 2 h. All absorbances were measured in a quartz cell at 625 nm in a UV-visible spectrophotometer (Thermo spectronic, Genesys 5, New York, EUA). The bacterial growth was estimated in CFU mL⁻¹ employing an absorbance adjustment with the McFarland 0.5 scale, as follows: CFU mL⁻¹ = (absorbance of medium with extract inoculated) (1.5 × 10⁸) / absorbance of the 0.5 McFarland scale.

Statistical analysis

All the experiments were conducted in triplicate (*n* = 3) and results were expressed as a mean ± standard deviation. The digestibility and prebiotic effect data were analyzed by ANOVA and Tukey mean comparison test (*P* ≤ 0.05). Graph Pad Prism-7[®] software was used for the statistical analysis.

RESULTS AND DISCUSSION

Plant material and extraction

There was no previous information on the yield percentages of the *S. queretaroensis* fruit, but in this study, a *Y* (%) of 91.23 ± 0.1 % was estimated. The yield of *S. queretaroensis* was similar to ultrasonic extraction from the pulp of *Hylocereus polyrhizus* (Cactaceae) of 90.08 % (Ramli *et al.*, 2014); therefore, the extraction yield

percentage depends on the conditions used for each extraction; however, the main advantage of the present procedure is that compounds of *S. queretaroensis* maintain their integrity, as the conditions used do not modify their chemical structure (Sasidharan *et al.*, 2011).

Physicochemical and proximal analyses

The *S. queretaroensis* extract from the red fruit possesses an acidic pH of 3.78 (Table 1), similar to that of other types of *S. queretaroensis* (pH 3.9 - 5.2) such as yellow fruit (Gaytán-Andrade *et al.*, 2019). Furthermore, the extracts from *S. griseus* and *Opuntia ficus-indica* have pH values of 4.2 and 6.04 ± 0.0 (Campos-Rojas *et al.*, 2011; Ochoa and Guerrero, 2012). It is clear that although all these extracts are acids, variation exists in acidic values among them. The difference in pH could be due to the concentration of chemical constituents (Harris, 2016). Furthermore, the pH can vary depending on the phenological conditions of the plant and the maturity of the fruit because its composition is related to the conditions of harvesting, post-harvest storage, biosynthesis, geographical region, season and age of the plant (Hernández *et al.*, 2017).

The extract of the red fruit had a lower TSS content (°Brix) (Table 1) than the fruit of the yellow one, which has 16 °Brix; however, its content is similar to that of *Hylocereus* spp. juice that contains 10-12 °Brix (Chuah *et al.*, 2008). The extract from *S. queretaroensis* fruit presents lower titratable acidity percentage (Table 1) than other fruits, including some species of *Stenocereus*. Since other types of *S. queretaroensis* have percentages of 0.15 to 0.50 % (Gaytán-Andrade *et al.*, 2019) and fruits of *Stenocereus* spp. from 0.25 to 0.28 % (Campos-Rojas *et al.*, 2011).

The production of citric acid corresponding to titratable acidity is due to the metabolic pathway of the plant and its phenological condition; therefore, its content varies according to the degree of maturity of the fruit. Furthermore, the citric acid content is a better indicator of flavor than pH (Sadler and Murphy, 2010); hence, the content detected

in the *S. queretaroensis* fruit extract confers it a pleasant taste due to its low acidity. TSS and titratable acidity play a role in the final flavor and quality of the extract; for this reason, they are relevant parameters in the food industry, thus, they are used as a base in the formulation of products (Campos-Rojas *et al.*, 2011).

The *S. queretaroensis* fruit extract displayed a reddish color and less luminosity (Table 1) than the *Hylocereus* spp. fruit pulp ($L^* = 51.35 \pm 0.7$, $a^* = 74.57 \pm 0.4$, $b^* = -1.34 \pm 0.5$) (Ochoa-Velasco *et al.*, 2012). The luminosity index makes the *S. queretaroensis* fruit extract more attractive, and the color is associated with the composition and maturity index of each fruit (Ortiz and Takahashi, 2015). The parameters a^* and b^* of *S. queretaroensis* slightly point out toward red and yellow, respectively. This coloration is linked to the content of betalains in some Cactaceae (Monroy-Gutiérrez *et al.*, 2017). At present, the color parameters of the extract of *S. queretaroensis* had not been described, but other studies showed that the whole fruit has $L^* 24.44 \pm 0.6$ to 34.44 ± 0.9 , $a^* 5.82 \pm 1.3$ to 18.82 ± 0.5 and $b^* 8 \pm 0.035$ to 16.75 ± 1.2 , which also correspond to shades of red and slight yellow (Soto-Castro *et al.*, 2019).

On the other hand, the moisture of the lyophilized extract from the *S. queretaroensis* fruit was lower (Table 1) than that of the extract of *S. queretaroensis* fruits and other non-lyophilized Cactaceae fruits. Moisture of 97.52 ± 0.1 to 98.88 ± 0.0 % has been determined in *S. queretaroensis* (Soto-Castro *et al.*, 2019), while is was 87.30 to 89.05 % in *Opuntia joconostle* (Contreras *et al.*, 2011), 78.4 to 88.27 % in *Opuntia ficus-indica* and 81.68 % in *Opuntia dillenii* (Díaz *et al.*, 2007). Nevertheless, lyophilization is a necessary step to standardize the extract on dry weight and to determine its retained moisture to establish the formulation of new products. Furthermore, knowing the moisture content allows establishing storage conditions to prevent microbiological damage (Qadri *et al.*, 2015).

The proximal analysis showed that the lyophilized extract of *S. queretaroensis* fruit does not contain minerals (Table

Table 1. Physicochemical and proximal analysis of the *S. queretaroensis* fruit extract.

Parameter	Value	Parameter	Value (%) w/w
pH	3.78 ± 0.1	Moisture	9.50 ± 0.7
Total soluble solids (° Brix)	11.30 ± 0.6	Ash	0.00 ± 0.0
Titratable acidity (citric acid %)	0.03 ± 0.0	Crude fat	0.00 ± 0.0
Color		Crude fiber	0.77 ± 0.0
L* (Lightness)	17.21 ± 0.1	Crude protein	1.36 ± 0.0
a* (+ red, - green)	38.61 ± 0.13	Nitrogen-free extract	88.82 ± 0.8
b* (+ yellow, - blue)	10.60 ± 0.23		

Values are means ± standard deviation, n = 3.

1); however, the ashes of the fruit could be retained in the pulp during the filtering process since the ash content is found at considerable levels in the fruits of other Cactaceae; for instance, 6.67 % in *S. griseus* (Gaytán-Andrade *et al.*, 2019) and 0.49 to 0.65 % in *O. joconostle* (Contreras *et al.*, 2011). The fat analysis showed that the lyophilized extract of *S. queretaroensis* does not contain fat (Table 1). The fruits of other Cactaceae have low-fat contents, thus, the fat from the lyophilized extract could be retained in the fruit pulp during filtration; *Opuntia dillenii* contains 0.71 % (Díaz *et al.*, 2007) and red *O. ficus-indica* contains less than 0.10 % fat (Repo de Carrasco and Encina, 2008). The lyophilized extract of *S. queretaroensis* showed a content of raw fiber (Table 1) lower than that of whole fruits from other Cactaceae. Because most of the fiber was retained during filtration, fiber content detected corresponds to the soluble fiber. The fruits of other Cactaceae have higher fiber contents; for example, *O. joconostle* contains from 2.35 to 4.28 % and *O. ficus-indica* from 0.38 to 3.9 % (Contreras *et al.*, 2011). The fiber content detected in the freeze-dried extract of *S. queretaroensis* fruit has the potential to improve gastrointestinal function, as fiber improves the digestion of nutrients and chemicals.

The lyophilized extract of *S. queretaroensis* fruit showed a protein content higher or similar to that of other Cactaceae fruits; therefore, it represents a source of nutrients. *Opuntia joconostle* showed a protein content of 0.71 to 1.56 % (Contreras *et al.*, 2011), *O. ficus-indica* of 0.9 to 1.5 % (Díaz *et al.*, 2007; Repo de Carrasco and Encina, 2008) and *O. dillenii* of 0.52 ± 0.12 . In addition, the *S. queretaroensis* extract showed a higher content of nitrogen-free extract (Table 1) than other Cactaceae as *O. joconostle* that contains 5.71–7.88 %, without considering the carbohydrate content, which is from 5.81 to 7.98 % (Contreras *et al.*, 2011), while *O. ficus-indica* (red variety) contains 16.7 % of the nitrogen-free extract, without including the carbohydrate content, corresponding to 20.2 % (Repo de Carrasco and Encina, 2008).

The fruits of different crops of *S. queretaroensis* contain 10–11 % of total sugars, but total solids, suspended solids, and vitamin C have not been detected (Gaytán-Andrade *et al.*, 2019). *Stenocereus griseus* fruit contains 9.9 % of total sugars (Chuck-Hernández *et al.*, 2016); consequently, it is believed that a large proportion of the nitrogen-free fraction of the lyophilized extract of *S. queretaroensis* corresponds to carbohydrates, vitamins and other compounds not determined in this study.

Digestibility

Digestibility tests of edible plant extracts are scarce because the methods for their *in vitro* evaluation are specific

(Bornhorst and Singh, 2014; Brodkorb *et al.*, 2019). In this context, no methods have been described to evaluate the digestibility of *S. queretaroensis*; nevertheless, simulated tests of oral *in vitro* digestibility (using saliva), gastric, and intestinal (small intestine and large intestine) are used for this purpose (Chen *et al.*, 2018); hence, this study focused on establishing the conditions for enzyme consumption (amylase), solubility (in gastric fluid simulated with pepsin), and microbial consumption of the *S. queretaroensis* extract (Chen *et al.*, 2018; Mkandawire *et al.*, 2015). By this way, we could infer the content of nutrients in the residue that remains from each digestive process (Gargallo *et al.*, 2006, Giraldo *et al.*, 2007) by using simulated processes where oral, gastric, and intestinal (large intestine) digestibility were measured based on the weight of the lyophilized extract consumed. The extract had an oral digestibility proportion of 89.42 ± 0.1 %, while the gastric showed 72.80 ± 4.2 % and the intestinal 97.45 ± 1.1 % (Figure 1); that is, the digestibility of the extract is different because the release of nutrients is different in each digestive process, and depends on the nature of the food matrix (Guerra *et al.*, 2012), as well as the mechanisms of degradation involved in each digestive process (Boland, 2016).

Currently, no oral digestibility tests have been described for this fruit or any similar one; however, results show that the amylase in the saliva consumes the carbohydrates contained in the extract since, according to the proximal analysis, it contains 88.82 ± 0.8 % of the nitrogen-free extract, which in this case corresponds mainly to sugars. The components of the extract are dissolved or hydrolyzed by salivary amylase and turn into smaller digestible carbohydrates (Boland, 2016), which could be studied. Additionally, there are many aspects of extract digestion that must be addressed to properly understand how the components of this food interact when ingesting it, to create new foods in the future (Boland, 2016). The gastric digestibility of the extract was performed by combining two methods; the procedure was established according to the composition of the extract as determined in the proximal analysis since no digestibility methods have been described for this plant.

In the assay, pepsin was used to hydrolyze the proteins in the extract and to measure the total content of insoluble compounds that persist as the remnant of digestion. Although the main content of the extract, according to the proximal analysis, corresponds to carbohydrates calculated as the nitrogen-free extract, the residue could contain non-digestible carbohydrates, crude fiber, insoluble phytochemical compounds and amino acids from protein hydrolysis (Gargallo *et al.*, 2006, Giraldo *et al.*, 2007). Also, the gastric fluid contains pepsin that solubilizes its main component (carbohydrates). As a result, high digestibility




Figure 1. Fractions of the oral, gastric, and intestinal digestibility of the fruit extract of *S. queretaroensis*. Data represent the mean \pm standard deviation ($n = 3$). Bars with the same letters show non-significant difference between each other (Tukey, $P \leq 0.05$).

values are obtained. The large intestine digestibility was adapted for humans by combining the Daisy^{II} and Chen methods (Chen *et al.*, 2018; Coblenz *et al.*, 2019). Until now, only the Daisy^{II} method has been described for ruminant digestibility. This method is efficient, fast, economical, and allows the simultaneous incubation of different samples in a jar (Giraldo *et al.*, 2007).

Results revealed that the percentage of intestinal digestibility is larger than the sum of the content of crude fiber, crude protein and nitrogen-free extract (90.95 ± 0.8 %) because the intestinal microbiota has the enzymatic capacity to degrade these components and the phytochemical compounds in the extract, which would correspond to around 6.50 ± 1.4 %. Therefore, intestinal digestibility was estimated as the extract digested by the human microbiota of the fecal inoculum (Gargallo *et al.*, 2006). As in the digestive process, the nutrients reach the intestinal microbiota to favor their growth and metabolite production (Ercolini and Fogliano, 2018); consequently, the microbiota protects the host from different diseases (Avolio *et al.*, 2020).

Prebiotic evaluation

The growth of *B. coagulans*, *L. sporogenes* and *L. delbrueckii* was higher in the medium supplemented with the fruit extract of *S. queretaroensis*, where they had $10.88 \times 10^{10} \pm 0.0$, $10.83 \times 10^{10} \pm 0.0$ and $10.46 \times 10^{10} \pm 0.0$ CFU mL⁻¹ at 24 h, while the growth of these microorganisms on MRS medium was $9.97 \times 10^{10} \pm 0.0$, $10.36 \times 10^{10} \pm 0.0$, and $9.38 \times 10^{10} \pm 0.0$ CFU mL⁻¹, respectively.

The extract did not modify the growth of *B. clausii* and *L.*

fermentum; these strains presented an exponential phase from 0 to 10 h, and then entered the stationary phase (Figure 2). During the exponential phase, MRS alone or added with extract contains the nutrients necessary for bacterial growth, which leads to cell division at a constant rate; however, when the microbial culture is consumed, it enters the stationary phase, so the microorganisms slow down their replication and die (Rolfe *et al.*, 2011). Nevertheless, some strains have different growth requirements and have enzymatic mechanisms (Boll *et al.*, 2014) to metabolize other carbon sources, such as the *S. queretaroensis* extract.

Furthermore, the bacterial growth depends not only on nutrients but also on factors such as enzymes, temperature and incubation periods (Dalié *et al.*, 2010). *B. coagulans* and *L. sporogenes* presented an exponential phase from 0 to 10 h, both in enriched medium and with MRS. *L. delbrueckii* presented an exponential phase from 0 to 12 h with the enriched medium and from 0 to 2 h with MRS; therefore, this strain accelerated growth because it quickly consumes nutrients. Accordingly, the enriched medium prolonged its exponential phase, as it provides an additional carbon source; however, *B. coagulans* and *L. sporogenes* (10 h) grew more in the enriched-medium than in MRS. These results suggest that extract from *S. queretaroensis* shows a prebiotic effect on *B. coagulans*, *L. sporogenes* and *L. delbrueckii*.

Bacillus coagulans and *L. esporogenes* showed growths of $10.67 \times 10^{10} \pm 0.0$ and $10.49 \times 10^{10} \pm 0.0$ CFU mL⁻¹ with the enriched medium, and of $9.75 \times 10^{10} \pm 0.0$ and $9.76 \times 10^{10} \pm 0.0$ CFU mL⁻¹ with MRS. Additionally, *L. delbrueckii* presented a growth of $10.42 \times 10^{10} \pm 0.0$ CFU mL⁻¹ (12 h) with the enriched medium and $8.40 \times 10^{10} \pm 0.0$ CFU mL⁻¹




Figure 2. Kinetics of *B. coagulans* (A), *L. sporogenes* (B), *B. clausii* (C), *L. fermentum* (D), and *L. delbrueckii* (E) growth in MRS medium enriched with extract of *S. queretaroensis* and MRS medium as control.

with the MRS (2 h). These biomasses are superior to that of *Lactobacillus casei* produced in the MRS medium (7.3×10^8 CFU mL^{-1}) (Jurado-Gómez *et al.*, 2014); thus, growth depends on the nutritional requirements of each strain. *Stenocereus queretaroensis* extract possesses prebiotic properties and high potential to be used as food matrix to develop new functional foods.

CONCLUSIONS

According to the results, the method developed for the extract from the *S. queretaroensis* fruit showed higher yields than other methods reported for cacti. The extract has an acidic pH, titratable acidity, TSS, and color that make it pleasant for consumption. Lyophilized drying removes moisture, thus helping to prevent deterioration,

very common in the fresh fruit of *S. queretaroensis*. The nutritional contribution of this fruit focuses on soluble crude fiber, protein content, and low nitrogen extract, which is related to high carbohydrate content; however, to strengthen knowledge about its nutritional value, it is necessary to deepen studies that include the profile of amino acids, vitamins, minerals and energy content. Furthermore, its nutritional composition is highly digestible at the oral, gastric and intestinal levels and gives it a prebiotic effect because its main component is a nitrogen-free extract, which is related to the content of highly soluble or degradable carbohydrates. Also, the developed methods could be useful to evaluate primary and secondary metabolites of the extract that are of interest to the food industry.

ACKNOWLEDGMENT

The authors thank CONACyT for the student scholarship (345654) and to Dr. José Angel Villarreal-Quintanilla and Dr. Eduardo García Martínez of the Antonio Narro Autonomous Agrarian University for identifying the plant and facilitating the development of the digestibility test, respectively.

BIBLIOGRAPHY

- AOAC, Association of Official Analytical Chemists (2000) Official Methods of Analysis of the Association of Official Analytical Chemists. Methods 925.10, 65.17, 974.24, 992.16. 17th edition. The Association of Official Analytical Chemists. Gaithersburg, Maryland, USA.
- Avolio E., P. Gualtieri, L. Romano, C. Pecorella, S. Ferraro, G. Palma, ... and A. De Lorenzo (2020) Obesity and body composition in man and woman: Associated diseases and the new role of gut microbiota. *Current Medicinal Chemistry* 27:216-229, <https://doi.org/10.2174/0929867326666190326113607>
- Boland M. (2016) Human digestion – a processing perspective. *Journal of the Science of Food and Agriculture* 96:2275-2283, <https://doi.org/10.1002/jsfa.7601>
- Boll M., C. Löffler, B. E. Morris and J. W. Kung (2014) Anaerobic degradation of homocyclic aromatic compounds via arylcarboxyl-coenzyme A esters: organisms, strategies and key enzymes. *Environmental Microbiology* 16:612-627, <https://doi.org/10.1111/1462-2920.12328>
- Bornhorst G. M. and R. P. Singh (2014) Gastric digestion *in vivo* and *in vitro*: how the structural aspects of food influence the digestion process. *Annual Review of Food Science and Technology* 5:111-132, <https://doi.org/10.1146/annurev-food-030713-092346>
- Brodtkorb A., L. Egger, M. Alming, P. Alvito, R. Assunção, S. Ballance, ... and I. Recio (2019) INFOGEST static *in vitro* simulation of gastrointestinal food digestion. *Nature Protocols* 14:991-1014, <https://doi.org/10.1038/s41596-018-0119-1>
- Campos-Rojas E., J. M. Pinedo-Espinoza, R. G. Campos-Montiel and A. D. Hernández-Fuentes (2011) Evaluation of pitaya plants (*Stenocereus* spp.) of natural populations of Monte Escobedo, Zacatecas. *Revista Chapingo Serie Horticultura* 17:173-182, <https://doi.org/10.5154/r.chsh.2011.17.028>
- Chen G., M. Xie, P. Wan, D. Chen, H. Ye, L. Chen, ... and Z. Liu (2018) Digestion under saliva, simulated gastric and small intestinal conditions and fermentation *in vitro* by human intestinal microbiota of polysaccharides from Fuzhuan brick tea. *Food Chemistry* 244:331-339, <https://doi.org/10.1016/j.foodchem.2017.10.074>
- Chuah T. G., H. L. Ling, N. L. Chin, T. S. Y. Choong and A. Fakhru'l-Razi (2008) Effects of temperatures on rheological behavior of dragon fruit (*Hylocereus* sp.) juice. *International Journal of Food Engineering* 4:1-28, <https://doi.org/10.2202/1556-3758.1519>
- Chuck-Hernández C., R. Parra-Saldivar and L. Sandate-Flores (2016) Pitaya (*Stenocereus* spp.). In: Encyclopedia of Food and Health. B. Caballero, P. Finglas and F. Toldrá (eds.). Elsevier Science and Technology. Boston, Massachusetts, USA. pp:385-391.
- Coblentz W. K., M. S. Akins, R. K. Ogden, L. M. Bauman and A. J. Stammer (2019) Effects of sample size on neutral detergent fiber digestibility of triticale forages using the Ankorm Daisy[®] Incubator system. *Journal of Dairy Science* 102:6987-6999, <https://doi.org/10.3168/jds.2019-16681>
- Contreras L. E., O. J. Jaimez, O. A. Castañeda, M. J. Añorve and R. S. Villanueva (2011) Sensory profile and chemical composition of *Opuntia joconostle* from Hidalgo, Mexico. *Journal of Stored Products and Postharvest Research* 2:37-39.
- Dalié D. K. D., A. M. Deschamps and F. Richard-Forget (2010) Lactic acid bacteria – Potential for control of mould growth and mycotoxins: a review. *Food Control* 21:370-380, <https://doi.org/10.1016/j.foodcont.2009.07.011>
- De U. C., S. Debnath, R. Ghosh and S. Debnath (2014) Preliminary screening for *in vitro* anti-enteric properties of a traditional herb *Dillenia pentagyna* Roxb. fruit extracts. *Asian Pacific Journal of Tropical Medicine* 7:S332-S341, [https://doi.org/10.1016/S1995-7645\(14\)60255-8](https://doi.org/10.1016/S1995-7645(14)60255-8)
- Díaz M. E. M., E. M. Rodríguez R. and C. Díaz R. (2007) Chemical characterization of *Opuntia dillenii* and *Opuntia ficus indica* fruits. *Food Chemistry* 103:38-45, <https://doi.org/10.1016/j.foodchem.2006.06.064>
- Dick M., L. Dal Magro, R. Costa R., A. de Oliveira R. and S. Hickmann F. (2019) Valorization of *Opuntia monacantha* (Willd.) Haw. cladodes to obtain a mucilage with hydrocolloid features: physicochemical and functional performance. *Macromolecules* 123:900-909, <https://doi.org/10.1016/j.jbiomac.2018.11.126>
- Ercolini D. and V. Fogliano (2018) Food design to feed the human gut microbiota. *Journal of Agricultural and Food Chemistry* 66:3754-3758, <https://doi.org/10.1021/acs.jafc.8b00456>
- Gargallo S., S. Calsamiglia and A. Ferret (2006) Technical note: A modified three-step *in vitro* procedure to determine intestinal digestion of proteins. *Journal of Animal Science* 84:2163-2167, <https://doi.org/10.2527/jas.2004-704>
- Gaytán-Andrade J. J., C. N. Aguilar-González, L. I. López-López, L. E. Cobos-Puc and S. Y. Silva-Belmares (2019) Perspectives for food development from pitayo *Stenocereus queretaroensis* (Weber) Buxbaum. In: Handbook of Research on Food Science and Technology. Vol. 1: Food Technology and Chemistry. M. L. Chávez-González, J. J. Buenrostro-Figueroa and C. N. Aguilar (eds.). Apple Academic Press. New Jersey, USA. pp:149-157.
- Giraldo L. A., L. A. Gutiérrez y C. Rúa (2007) Comparación de dos técnicas *in vitro* e *in situ* para estimar la digestibilidad verdadera en varios forrajes tropicales. *Revista Colombiana de Ciencias Pecuarias* 20:269-279.
- Guerra A., L. Etienne-Mesmin, V. Livrelli, S. Denis, S. Blanquet-Diot and M. Alric (2012) Relevance and challenges in modeling human gastric and small intestinal digestion. *Trends in Biotechnology* 30:591-600, <https://doi.org/10.1016/j.tibtech.2012.08.001>
- Harris D. C. (2016) Análisis Químico Cuantitativo. Tercera edición. Editorial Reverté. Barcelona, España. 894 p.
- Hernández O. L., P. Carranza R., L. E. Cobos P., L. I. López L., J. A. Ascasio V. and S. Y. Silva B. (2017) Bioguided fractionation from *Solanum elaeagnifolium* to evaluate toxicity on cellular lines and breast tumor explants. *Vitae* 24:124-131, <https://doi.org/10.17533/udea.vitae.v24n2a05>
- Jurado-Gómez H., F. Calpa-Yamá y A. Chaspuengal-Tulcán (2014) Determinación de parámetros cinéticos de *Lactobacillus casei* en dos medios probióticos. *Veterinaria y Zootecnia* 8:15-35, <https://doi.org/10.17151/vetzo.2014.8.2.2>
- Mkandawire N. L., S. A. Weier, C. L. Weller, D. S. Jackson and D. J. Rose (2015) Composition, *in vitro* digestibility, and sensory evaluation of extruded whole grain sorghum breakfast cereals. *LWT - Food Science and Technology* 62:662-667, <https://doi.org/10.1016/j.lwt.2014.12.051>
- Monroy-Gutiérrez T., M. T. Martínez-Damián, A. F. Barrientos-Priego, C. Gallegos-Vázquez, O. Cruz-Alvarez y H. Vargas-Madriz (2017) Compuestos bioactivos y capacidad antioxidante en frutos de xocotuna, tuna y xoconostle (*Opuntia* spp.). *Chilean*

- Journal of Agricultural & Animal Sciences* 33:263-272, <https://doi.org/10.4067/S0719-38902017005000704>
- Ochoa C. E. y J. A. Guerrero (2012) Efecto del almacenamiento a diferentes temperaturas sobre la calidad de tuna roja (*Opuntia ficus indica* (L.) Miller). *Información Tecnológica* 23:117-128, <https://doi.org/10.4067/S0718-07642012000100013>
- Ochoa-Velasco C. E., V. García-Vidal, J. J. Luna-Guevara, M. L. Luna-Guevara, P. Hernández-Carranza y J. A. Guerrero-Beltrán (2012) Características antioxidantes, fisicoquímicas y microbiológicas de jugo fermentado y sin fermentar de tres variedades de pitahaya (*Hylocereus* spp). *Scientia Agropecuaria* 3:279-289.
- Orozco-Sifuentes M. M., J. E. García-Martínez, C. A. Arévalo-Sanmiguel, F. Ramírez-Godina y M. H. Reyes-Valdés (2019) Nutritive potential of soto (*Dasyllirion cedrosanum*) seeds. *Revista Fitotecnia Mexicana* 42:385-392, <https://doi.org/10.35196/rfm.2019.4.385-392>
- Ortiz T. A. and L. S. A. Takahashi (2015) Physical and chemical characteristics of pitaya fruits at physiological maturity. *Genetics and Molecular Research* 14:14422-14439, <https://doi.org/10.4238/2015.November.18.5>
- Qadri O. S., B. Yousuf and A. K. Srivastava (2015) Fresh-cut fruits and vegetables: critical factors influencing microbiology and novel approaches to prevent microbial risks - A review. *Cogent Food & Agriculture* 1:1121606, <https://doi.org/10.1080/23311932.2015.1121606>
- Quiroz-González B., R. García-Mateos, J. J. E. Corrales-García and M. T. Colinas-León (2018) Pitaya (*Stenocereus* spp.): an under-utilized fruit. *Journal of the Professional Association for Cactus Development* 20:82-100.
- Ramírez-Jaspeado R., N. Palacios-Rojas, M. Nutti y S. Pérez (2020) Estados potenciales en México para la producción y consumo de frijol biofortificado con hierro y zinc. *Revista Fitotecnia Mexicana* 43:11-23, <https://doi.org/10.35196/rfm.2020.1.11>
- Ramli N. S., P. Ismail and A. Rahmat (2014) Influence of conventional and ultrasonic-assisted extraction on phenolic contents, betacyanin contents, and antioxidant capacity of red dragon fruit (*Hylocereus polyrhizus*). *The Scientific World Journal* 2014:964731, <https://doi.org/10.1155/2014/964731>
- Repo de Carrasco R. and C. R. Encina Z. (2008) Determinación de la capacidad antioxidante y compuestos bioactivos de frutas nativas peruanas. *Revista de la Sociedad Química del Perú* 74:108-124.
- Rolfé M. D., C. J. Rice, S. Lucchini, C. Pin, A. Thompson, A. D. S. Cameron, ... and J. C. D. Hinton (2011) Lag phase is a distinct growth phase that prepares bacteria for exponential growth and involves transient metal accumulation. *Journal of Bacteriology* 194:686-701, <https://doi.org/10.1128/JB.06112-11>
- Román-Cortés N. R., M. R. García-Mateos, A. M. Castillo-González, J. Sahagún-Castellanos y M. A. Jiménez-Arellanes (2018) Características nutricionales y nutraceuticas de hortalizas de uso ancestral en México. *Revista Fitotecnia Mexicana* 41:245-253, <https://doi.org/10.35196/rfm.2018.3.245-253>
- Sadler G. D. and P. A. Murphy (2010) pH and titratable acidity. In: Food Analysis. S. S. Nielsen (ed.). Springer. Boston, USA. pp:219-238, https://doi.org/10.1007/978-1-4419-1478-1_13
- Sasidharan S., Y. Chen, D. Saravanan, K. M. Sundram and L. Yoga Latha (2011) Extraction, isolation, and characterization of bioactive compounds from plants' extracts. *African Journal of Traditional, Complementary and Alternative Medicines* 8:1-10.
- SIAP, Servicio de Información Agroalimentaria y Pesquera (2019) Anuario estadístico de la producción Agrícola. Secretaría de Agricultura y Desarrollo Rural. Ciudad de México. <https://nube.siap.gob.mx/cierreagricola/> (Agosto 2019).
- Soto-Castro D., M. Chávez G., F. M. León-Martínez, P. A. Santiago-García, I. Aragón-Lucero I. and F. Antonio-Antonio (2019) Spray drying microencapsulation of betalain rich extracts from *Escontria chiotilla* and *Stenocereus queretaroensis* fruits using cactus mucilage. *Food Chemistry* 272:715-722, <https://doi.org/10.1016/j.foodchem.2018.08.069>
- Talebi S., M. Bagherniya, S. L. Atkin, G. Askari, H. M. Orafai and A. Sahebkar (2020) The beneficial effects of nutraceuticals and natural products on small dense LDL levels, LDL particle number and LDL particle size: a clinical review. *Lipids in Health and Disease* 19:66, <https://doi.org/10.1186/s12944-020-01250-6>
- Ye Z., C. Cao, Y. Liu, P. Cao and Q. Li (2018) Digestion fates of different edible oils vary with their composition specificities and interactions with bile salts. *Food Research International* 111:281-290, <https://doi.org/10.1016/j.foodres.2018.05.040>
- Yoon C., D. R. Jacobs, D. A. Duprez, D. Neumark-Sztainer, L. M. Steffen and S. M. Mason (2019) Problematic eating behaviors and attitudes predict long-term incident metabolic syndrome and diabetes: the coronary artery risk development in young adults study. *International Journal of Eating Disorders* 52:304-308, <https://doi.org/10.1002/eat.23020>

