



PROTEIN SOLUTIONS FROM *Xanthomonas* AS RESISTANCE INDUCTORS FOR THE CONTROL OF BACTERIAL SPOT IN TOMATO

SOLUCIONES PROTEICAS DE *Xanthomonas* COMO INDUCTORES DE RESISTENCIA PARA EL CONTROL DE LA MANCHA BACTERIANA EN JITOMATE

Sergio D. Valerio-Landa¹, Evangelina E. Quiñones-Aguilar¹, Jhony N. Enriquez-Vara¹, Rodolfo Hernández-Gutierrez², Luis G. Hernández-Montiel³ and Gabriel Rincon-Enriquez^{1*}

¹Centro de Investigación y Asistencia en Tecnología y Diseño del Estado de Jalisco A.C. (CIATEJ), Laboratorio de Fitopatología de Biotecnología Vegetal, Zapopan, Jalisco, México. ²CIATEJ, Biotecnología Médica y Farmacéutica, Guadalajara, Jalisco, México. ³Centro de Investigaciones Biológicas del Noroeste, Laboratorio de Fitopatología, La Paz, Baja California Sur, México.

*Corresponding author (grincon@ciatej.mx)

SUMMARY

Bacterial spot, caused by *Xanthomonas vesicatoria*, is a tomato (*Solanum lycopersicum*) disease and its control is usually with chemical products. However, there are alternatives to mitigate diseases, such as activation of resistance state in plants through pathogen-associated molecular patterns (PAMP) inductors. In this study the effect of protein solutions from *X. vesicatoria* were evaluated as PAMP inductors and activation of resistance state in tomato against bacterial spot under *in vitro* conditions. Protein solutions from *X. vesicatoria* strain BV801 (BV801) were obtained, and its protein profile was determined by SDS-PAGE. The BV801 protein solutions with antibacterial activity against *X. vesicatoria* was evaluated in culture medium with the double-layer plaque assays. The *in vitro* resistance induction trial in tomato was carried out by means of a completely randomized experimental design with 8 treatments: seven protein concentrations (0.0, 0.001, 0.01, 0.1, 1.0, 10 and 100 µg mL⁻¹) inoculated with BV801 and a control of healthy plants (without proteins and without BV801). The results showed that 19 major protein bands formed the protein profile of BV801, which did not show antibacterial activity *in vitro* against *X. vesicatoria*. A 50 % reduction in the disease caused by the bacterial spot was found in tomato plants treated with protein solutions from BV801, at concentrations 0.1, 1.0 and 10 µg mL⁻¹ (Kruskal-Wallis, $P \leq 0.05$); furthermore, plant growth maintained statistical similarity with healthy control plants (Tukey, Bonferroni, $P \leq 0.05$). The protein solutions from BV801 are PAMP inductors and increased the tomato plants resistance to bacterial spot disease.

Keywords: *Xanthomonas vesicatoria*, elicitor, ISR, resistance inductor, SAR.

RESUMEN

La mancha bacteriana causada por *Xanthomonas vesicatoria* es una enfermedad del jitomate (*Solanum lycopersicum*) y su control es generalmente con productos químicos. Sin embargo, existen alternativas para combatir este tipo de enfermedades, como la activación del estado de resistencia en plantas a través de inductores de patrones moleculares asociados a patógenos (PMAP). En este estudio se evaluó el efecto de soluciones proteicas de *X. vesicatoria* como inductores PMAP y activadores del estado de resistencia en jitomate contra la mancha bacteriana en condiciones *in vitro*. Se obtuvieron soluciones proteicas de *X. vesicatoria* cepa BV801 (BV801) y su perfil de proteínas se determinó mediante SDS-PAGE. Se evaluó la actividad antibacteriana de soluciones proteicas de BV801 contra *X. vesicatoria* en medio de cultivo con la técnica de doble placa. Para el ensayo *in vitro* de inducción de resistencia

en jitomate se utilizó un diseño experimental completamente al azar con 8 tratamientos: 7 concentraciones de proteínas (0.0, 0.001, 0.01, 0.1, 1.0, 10 y 100 µg mL⁻¹) inoculadas con BV801 y un control de plantas sanas (sin proteínas y sin BV801). Los resultados mostraron que 19 bandas de proteínas principales formaron el perfil proteico de BV801, las cuales no mostraron actividad antibacteriana *in vitro* contra *X. vesicatoria*. Se encontró una reducción del 50 % en la enfermedad causada por la mancha bacteriana en las plantas de jitomate tratadas con soluciones proteicas de BV801, en las concentraciones 0.1, 1.0 y 10 µg mL⁻¹ (Kruskal-Wallis, $P \leq 0.05$); además, el crecimiento vegetal mantuvo similitud estadística con plantas del control sano (Tukey, Bonferroni, $P \leq 0.05$). Las soluciones de proteínas de BV801 mostraron ser inductores PMAP y aumentaron la resistencia de las plantas de jitomate a la enfermedad de las manchas bacterianas.

Palabras clave: *Xanthomonas vesicatoria*, elicitor, inductor de resistencia, ISR, SAR.

INTRODUCTION

In Mexico, tomato cultivation is one of the most important agricultural activities, with a record production of 3,441,639.37 t in 2019 (SIAP, 2021). Various abiotic and biotic stresses affect tomato production and fruit quality. Among the biotic stresses, diseases are one of the most important challenging components, leading to serious effects on plant growth and productivity (Sharma and Bhattarai, 2019). The bacterial spot, caused by *Xanthomonas vesicatoria*, *X. euvesicatoria*, *X. perforans* and *X. gardneri*, is one of the most important diseases in tomato (Jones *et al.*, 2004; Strayer *et al.*, 2016). The plants with bacterial spot disease present small yellow-green lesions on leaves, progressing into light dark, and as the infection progresses the leaves become deformed and develop dark necrotic lesions (Vallad *et al.*, 2004). In addition, this infection leads to defoliation and fruits with lesion, ultimately causing severe reduction in tomato productivity (Potnis *et al.*, 2015). The control of bacterial spot in field is challenging, especially during warm and humid conditions, which characterize the main tomato

producing areas in Mexico.

The chemical control has been a primary focus in past management strategies for bacterial spot tomato. The application in plants of copper-based bactericides, alone or combined with fungicides, such as mancozeb (ethylene-bis-dithiocarbamates), and antibiotics, such as streptomycin or kasugamycin, are the main strategies for the control of bacterial spot (Vallad *et al.*, 2010; Šević *et al.*, 2019). However, races of *Xanthomonas* resistant to copper and streptomycin are commonly present in field since 1960s (Thayer and Stall, 1961). In addition, the genes that encode resistance to copper and streptomycin can be transmitted through plasmids, allowing the rapid spread of resistance to other non-resistant bacteria (Ritchie, 2000). On the other hand, breeding programs to confer resistance to tomato plants against bacterial spot by insertion of resistance genes (*R-genes*) have been unsuccessful, due to apparition of new strains of the pathogen that overcome resistance (Adhikari *et al.*, 2020).

Due to the high resistance of *Xanthomonas* to bactericides and the limited success of breeding programs to provided resistance to tomato plants against the bacterial spot, the identification and exploration for alternative control strategies become mandatory. The application of inducers in plants to confer a physiological state of resistance against phytopathogens has shown be alternative strategy in the control of diseases (Dewen *et al.*, 2017). In addition, the resistance state in tomato induced against the bacterial spot has shown levels of efficiency, similar or superior to the standard copper bactericide control methods (Louws *et al.*, 2001).

The plants have developed a unique ability to identify the pathogen microbes that cause infection through conserved microbial molecules known as pathogen-associated molecular pattern (PAMP) and these conserved molecules are elicitors of immunity system (Boller and Felix, 2009; Zipfel, 2014). On the other hand, extracellular immune receptor in the plant cell known as pattern recognition receptors (PRR) identify PAMP (Bhattarai *et al.*, 2016). The perception of PAMP by PRR triggers the signaling of a state of immunity in plants known as immunity triggered by PAMP (ITP). This state of immunity confers resistance at local sites of infection of pathogens and contributes to the signaling for the systemic activation of resistance against future pathogenic attacks (Wu *et al.*, 2014).

The use of PAMP inducers, e.g. proteins, peptides, oligosaccharides, lipids, nucleic acids, or any other cellular component, have shown relevance as an effective strategy for the control of phytopathogens (Dewen *et al.*, 2017). Particularly, proteins obtained from *Bacillus*

amyloliquefaciens and *B. pumilus* induce resistance against bacterial spot of *X. vesicatoria* in tomato (Lanna-Filho *et al.*, 2013); also, polysaccharides obtained from *X. gardneri* induce resistance to bacterial spot in tomato (Luiz *et al.*, 2015). In contrast, there are few reports about the ability of biological molecules belonging to *X. vesicatoria* to protect the plants against bacterial diseases (Gottig *et al.*, 2018). The aim of this study was to determinate the effect of *X. vesicatoria* protein solutions as inducers of resistance against bacterial spot in tomato plants under *in vitro* conditions.

MATERIALS AND METHODS

Phytopathogenic bacteria and plant material

The BV801 strain of *X. vesicatoria* (BV801) was provided by the Phytopathology Laboratory of CIATEJ and was cultivated in NYG liquid medium (0.5 % peptone (N), 0.3 % yeast extract (Y), 2 % glycerol (G)) at 28 °C for 36 h on a rotary shaker at 200 rpm. The plant material was obtained from tomato seeds cv. Maya (Seminis®), which were sterilized with 3 % sodium hypochlorite for 5 min and washed with sterile distilled water. The seeds were germinated *in vitro* on MS medium (Murashige and Skoog, 1962) at 26 °C for 5 d. The seedlings were transferred to Half-closed Arbuscular Mycorrhizal - Plant (HAM-P) system described by Voets *et al.* (2005), with the following modification: a 12 mm diameter perforation was made on the lid of the Petri dish. Through this hole, the MS medium was renewal each 7 days and a sterile rubber plug was used to cover the perforation (Figure 1).




Figure 1. Modified Half-closed Arbuscular Mycorrhizal-Plant (HAM-P) system (Voets *et al.*, 2005) with modifications (rubber stopper).

Protein extraction and quantification

The BV801 strain cultivated in NYG liquid medium was collected by centrifugation at 4 °C for 10 min at 8,000 rpm, and was washed three times with 1X PBS (Phosphate buffered Saline). The cells were lysed in SDS-Tris buffer (0.1 % Sodium Dodecyl Sulfate, Tris-base 100 mM and adjusted to pH 9.5), and the cell disruption was performed in a ultrasonicator at 20 kHz at 4 °C for 2 min with intervals of 30 s. The sample was centrifuged at 4 °C for 30 min and 12,000 rpm and the supernatants were used for determination of soluble proteins. Protein quantification was performed using the Bradford assay and the BSA protein standard (Bradford, 1976). To determine the protein profile, the protein solutions was adjusted to a concentration of 31.25, 62.5, 125, 250, 500 and 1000 µg mL⁻¹. For the tests of antibacterial activity induction of plant resistance proteins, solutions were adjusted in the concentrations of 0.001, 0.01, 0.1, 1.0, 10 and 100 µg mL⁻¹. Protein solutions were stored at -20 °C until use.

Protein profile of *X. vesicatoria* strain BV801 by SDS-PAGE

The SDS-PAGE protocol (sodium dodecyl sulfate polyacrylamide gel electrophoresis) was used for protein profiling as described by Laemmli (1970). The stacking gel contained 4 % acrylamide (0.5 M Tris-HCl, adjusted to pH 6.8 10 % SDS w/v) and resolution of gel of acrylamide 12 % (1.5 M Tris-HCl, pH 8.8, 10 % SDS w/v) was made. Samples of 5 µL of protein solutions (31.25, 62.5, 125, 250, 500 and 1000 µg mL⁻¹) were loaded with 4.75 µL of Laemmli sample buffer and 0.25 µL of β-mercaptoethanol and denatured at 95 °C for 5 min. Electrophoresis was performed in 1X SDS-PAGE buffer (250 mM tris, 1.92 M glycine, 1 % SDS, pH 8.3). The electrophoresis conditions were 120 volts at 80 amps per 100 min. Gel staining with Coomassie R-250 (50 % methanol, 10 % acetic acid, 0.25 % coomassie blue R-250) was performed. The excessing dye solution was removed with fading solution. The visualization and capture of images of the gels were carried out in a photodocumentor (Bio-Rad®, Gel Doc XR System).

Antibacterial activity of protein solutions

Antibacterial activity of BV801 proteins solutions were determinate *in vitro* test of inhibition in double-layer plaque as described by Cavalcanti *et al.* (2006) with minor modifications. BV801 bacterial strain was cultivated in NYG liquid medium and concentration was adjusted at 1×10⁸ CFU mL⁻¹ using a spectrophotometer (Eppendorf®, BioPhotometer) at 600 nm and optical density (OD) of 1. The double-layer plaque was prepared by adding 100 µL of bacterial suspension OD = 1 to 4.9 mL of NYGA (0.5 %

peptone, 0.3 % yeast extract, 2 % glycerol, 0.7 % agar) at 40 °C and allowed to gel at room temperature. The different proteins solutions (0.001, 0.01, 0.1, 1.0, 10 and 100 µg mL⁻¹) and SDS-Tris solution (dissolution buffer of proteins without protein) were added in drops of 10 µL on the double-layer plaque and allowed to dry under hood flow. Neomycin (1 µg µL⁻¹) and sterile distilled water were used as control positive and negative of inhibitors, respectively. The Petri plates were incubated at 28 °C for 48 h and the inhibition halo of *X. vesicatoria* by proteins solutions and controls were registered. The experimental design was completely randomized and six repetitions per treatment were used.

Induction of resistance test in tomato against bacterial spot

An experiment was conducted in a completely randomized design with eight treatments and seven repetitions. The treatments were plants inoculated with the strain BV801 of *X. vesicatoria* and the application of 7 concentrations of *Xanthomonas* proteins solutions: 0.0 (diseased plants), 0.001, 0.01, 0.1, 1.0, 10 and 100 µg mL⁻¹, and a control without protein and without phytopathogenic bacteria (healthy plants). The protein solutions were sprayed on both sides (adaxial and abaxial) of leaves with the help of a nozzle sprayer (≈0.5 mL / plant) (ATO50 Truper®). The protein solutions were subjected to 95 °C for 5 min, to promote the deployment of secondary protein structure and promote epidermal exposure. Protein solutions are allowed to cool to room temperature before being applied to the plant. After 36 h, the bacterial (BV801 strain) suspension (1×10⁸ CFU mL⁻¹) was applied on leaves. The plants were kept at 28 °C, 55 % HR and 10/14 h light/dark in bioclimatic chamber (SEV-PRENDÓ-México, INCL-20) for 14 days.

Severity of disease in tomato plants and plant growth parameters

The severity of *X. vesicatoria* in tomato plants was determined with a scale of 0 to 5 (Figure 2). The scale was based on percentage of tissue damaged due to inoculation of *X. vesicatoria* (0 = healthy plant, 1 = 20 %, 2 = 40 %, 3 = 60 %, 4 = 80 % and 5 = dead plant). The plant height (cm), stem diameter (mm), root length (cm), fresh biomass (g), dry biomass (g), number of branches and number of leaves were quantified at the end of the experiment.

Statistical data analysis

The severity scale was analyzed using Kruskal-Wallis nonparametric technique and median confidence interval ($P \leq 0.05$). One-way analysis of variance (ANOVA) was



Figure 2. Severity scale of bacterial spot in tomato plants *in vitro* conditions by effect of inoculation of *X. vesicatoria*. The scale is based on percentage of tissue damaged due to inoculation of *X. vesicatoria*. 0: healthy plant, 1: 20 %, 2: 40 %, 3: 60 %, 4: 80 % and 5: dead plant.

performed on growth variables, that were compared with Tukey ($P \leq 0.05$). A Bonferroni statistical test was used for dry biomass. The data were analyzed using Statgraphics centurion XVII (<http://www.statgraphics.com/>).

RESULTS

Protein solutions profile of *X. vesicatoria* and antibacterial activity

The proteins profile on one dimension were obtained from *X. vesicatoria*, and approximately 35 bands conform the protein profile ($1000 \mu\text{g mL}^{-1}$). Particularly, 19 protein bands have a higher proportion compared to the 16 others protein bands, evidenced by the dilution of 125 and $250 \mu\text{g mL}^{-1}$ (Figure 3). In the image, the arrows on the right indicate the position of the 19 main protein bands, whose estimated molecular weights were 10, 12, 15, 17, 20, 23, 25, 30, 35, 40, 45, 55, 60, 65, 70, 80, 100, 110, and 200 kDa. The bacterial growth inhibition test shown that only the antibiotic neomycin ($1 \mu\text{g } \mu\text{L}^{-1}$) had a direct inhibitory effect on the growth of *X. vesicatoria* (Figure 4). These results indicate the absence of compounds with antibacterial activity in the protein solutions from strain BV801 and in the protein dissolution buffer (SDS-Tris) solution.

Effects of protein solutions on the disease severity of bacterial spot and growth plant

Significant effects were found in the reduction of the severity of the bacterial spot (95 % median confidence interval), due to the application of protein solutions from *X. vesicatoria*. Particularly, plants treated with protein

solutions 0.1 , 1.0 and $10 \mu\text{g mL}^{-1}$ showed a significant reduction on the severity scale by 55, 50 and 50 %, respectively, with respect to diseased plants (without protein solutions and inoculated with *X. vesicatoria*) (Figure 5). The protective effect of these protein solution treatments was demonstrated by significant reduction (test Kruskal-Wallis, $P = 0.0035$) in the severity of disease at 14 days after exposure to phytopathogenic bacteria (Figure 6). On the other hand, the biomass of plants (wet and dry) treated with protein solutions and healthy plants (without protein solutions and without *X. vesicatoria*) were statistically equal, but diseased plants (without protein solutions and inoculated with BV801) were significantly (Tukey and Bonferroni, $P \leq 0.05$) lesser than healthy plants (Table 1). Likewise, the number of leaves and root length of healthy plants were significantly (Tukey, $P \leq 0.05$) higher to diseased plants untreated with protein solutions. On the other hand, all growth variables of plants treated with 0.1 and $1.0 \mu\text{g mL}^{-1}$ protein solutions were statistically equal (Tukey, $P \leq 0.05$) to healthy plants. No statistical differences were observed in plant height, stem diameter and number of branches.

DISCUSSION

The use of PAMP inductors in tomato to activate resistance state against phytopathogens has been an effective strategy in the control of bacterial spot infection (Blainski *et al.*, 2017; Luiz *et al.*, 2015; Lanna-filho *et al.*, 2013; Cavalcanti *et al.*, 2007). According to Cavalcanti *et al.* (2007), a state of resistance induced in plants is characterized by the reduction of the size and number of injuries that develop after the inoculation of a phytopathogen, which agrees with the reduction of scale




Figure 3. Protein profile obtained from *X. vesicatoria* visualized by SDS-PAGE method. The concentration of protein loaded in each lane is indicated at the top of the image. Protein ladder molecular weight of 10 to 250 kDa (PageRuler™ Plus Prestained Protein Ladder, Thermo Scientific™). The arrows on the right indicate the position of the 19 main protein bands.




Figure 4. Bacterial growth inhibition test. The plates show experimental replicates. The numbers on the plates indicate the concentration of protein solutions applied in $\mu\text{g mL}^{-1}$. SDS: protein dissolution buffer, H_2O : sterile distilled water (negative inhibition control), antibiotic neomycin (positive inhibition control: 10 μL at 1 $\mu\text{g mL}^{-1}$).

of severity in plants observed when applying treatments of 0.1, 1.0 and 10 $\mu\text{g mL}^{-1}$ of proteins solutions. Particularly, protein solutions from BV801 showed no capacity to inhibit *in vitro* growth of the *X. vesicatoria*, which suggests the absence of compounds with antibacterial activity in these protein solutions. Therefore, in this work the control mechanisms of the bacterial spot in tomato observed were related to the induction of resistance through PAMP inductors.

Particularly, the use of resistance inductors in tomato has shown the activation of resistance mechanisms that

activate enzymes involved in resistance to oxidative stress such as catalase, superoxide dismutase and glutathione reductase (Blainski *et al.*, 2018), and increased gene expression of pathogenesis-related proteins (PR-genes), PR1a and PR1b (Huang and Vallad, 2018). Systemic acquired resistance (SAR) and induced systemic resistance (ISR) are two forms of induced resistance, wherein plant defences are preconditioned by prior infection or treatment, that results in resistance against subsequent challenge by a pathogen (Reimer-Michalski and Conrath, 2016). The SAR is characterized by pathogen-related proteins (PR proteins) and can enhance cellular responses and




Figure 5. Effect of the application of bacterial protein solutions in tomato plants against bacterial spot caused by *X. vesicatoria* under *in vitro* conditions. Different letters indicate statistical differences according to the nonparametric test of Kruskal-Wallis ($P = 0.0035$) and 95 % confidence intervals of median. The bar represents the standard error.




Figure 6. Plants treatments with *X. vesicatoria* strain BV801 plus protein solutions ($\mu\text{g mL}^{-1}$): a) 0.001, b) 0.01, c) 0.1, d) 1.0, e) 10, f) 100, g) Diseased plants (without protein solutions and inoculated with *X. vesicatoria*) and h) Healthy plants (without protein solutions and without *X. vesicatoria*). Left side: scale with length = 14 cm.

Table 1. Effect of application of bacterial protein solutions on plant growth of tomato plants under *in vitro* conditions.

Treatment		Plant height (cm)	Stem diameter (mm)	Number of branches	Number of leaf	Root length (cm)	Total biomass (g)	
Strain BV801	Protein solutions ($\mu\text{g mL}^{-1}$)						Wet	Dry*
With	100	5.9±0.9 a	1.7±0.2 a	4.2±0.5 a	11.8±1.6 ab	12.7±2.8 ab	0.9±0.3 ab	0.05±0.1 b
	10	6.3±0.2 a	1.8±0.1 a	3.7±0.2 a	10.2±1.0 b	11.0±2.1 ab	0.8±0.2 ab	0.07±0.1 ab
	1.0	6.4±0.7 a	1.8±0.2 a	4.3±0.3 a	11.8±1.7 ab	13.7±2.6 ab	0.9±0.2 ab	0.06±0.1 ab
	0.1	6.4±0.3 a	1.9±0.1 a	4.3±0.2 a	11.7±1.0 ab	16.0±1.3 a	0.9±0.1 ab	0.07±0.1 ab
	0.01	5.7±0.3 a	1.9±0.1 a	3.7±0.2 a	10.5±0.6 ab	11.3±0.9 ab	0.7±0.1 b	0.05±0.1 b
	0.001	5.9±0.5 a	1.8±0.1 a	3.7±0.2 a	9.30±0.7 b	11.8±1.4 ab	0.7±0.2 ab	0.04±0.1 b
	0†	5.3±0.9 a	1.5±0.2 a	3.0±0.4 a	7.00±0.9 b	8.3±1.00 b	0.6±0.1 b	0.04±0.1 b
Without	0‡	6.7±0.5 a	2.1±0.1 a	4.1±0.3 a	15.7±1.5 a	16.4±1.5 a	1.4±0.2 a	0.10±0.1 a

Means with the same letter in column indicate no statistical differences according to Tukey and *Bonferroni statistical test ($P \leq 0.05$). Standard error (\pm). †Diseased plants: without protein solutions and inoculated with BV801. ‡Healthy plants: without protein solutions and without BV801.

defense against phytopathogens. In *Arabidopsis*, localized applications PAMP showed elevated levels of salicylic acid, the SAR regulatory metabolite and PR-genes expression (Mishina and Zeier, 2007). Together, this mechanisms limit progress and subsequent development of phytopathogens attack. Probably, these mechanisms could have been present in tomato plants treated with protein solutions of 0.1, 1.0 and 10 $\mu\text{g mL}^{-1}$, treatments that showed less damage to infection by *X. vesicatoria*. However, further studies are necessary to corroborate these assumptions.

In regard to the use of PAMP inductors to activate resistance, in *Xanthomonas* genus has been reported that pathogenicity and hypersensitive response proteins (Hrp-proteins) of 20 to 25 kDa are PAMPs and that are involved in the induction of the defense response of plants (Gottig *et al.*, 2018). In this research, the resolution of the BV801 protein profile showed the accumulation of protein bands 20, 23 and 25 kDa, probably related with Hrp-proteins. On the other hand, it has been indicated that flagellin peptides of 22 amino acids from *Xanthomonas* (flg22-Xac) are PAMPs that induce resistance to bacterial spot in tomato, and whose immune perception is mediated by PRR as FLS2 described in *Arabidopsis* (Bhattarai *et al.*, 2016). Bacterial flagellin is related with the motility of bacterial and is a PAMP preserved in *Xanthomonas* and it could be present in the BV801 protein solutions evaluated with inductors. However, more studies are necessary to confirm that flagellin peptides, as well as Hrp-proteins, are present in protein solutions of BV801.

CONCLUSION

In this research, we demonstrate that *X. vesicatoria* strain BV801 protein solutions contain PAMP that maintain

eliciting activity of the plant immunity system and that they could be involved in the activation of the resistance state against pathogens. This protein solution did not show antibacterial activity. The presence of proteins that maintained PAMP activity allowed them to be recognized by the immune system of *S. lycopersicum*, leading to the activation of a state of resistance.

ACKNOWLEDGEMENTS

This research was supported by projects of the Plant Pathology Laboratory of CIATEJ: Problemas Nacionales clave CONACyT: 2015-01-338 and Laboratorio Nacional PLANTECC clave 315918-2021.

BIBLIOGRAPHY

- Adhikari P., T. B. Adhikari, F. J. Louws and D. R. Panthee (2020) Advances and challenges in bacterial spot resistance breeding in tomato (*Solanum lycopersicum* L.). *International Journal of Molecular Sciences* 21:1734, <https://doi.org/10.3390/ijms21051734>
- Bhattarai K., F. J. Louws, J. D. Williamson and D. R. Panthee (2016) Differential response of tomato genotypes to *Xanthomonas*-specific pathogen-associated molecular patterns and correlation with bacterial spot (*Xanthomonas perforans*) resistance. *Horticulture Research* 3:16035, <https://doi.org/10.1038/hortres.2016.35>
- Blainski J. M. L., A. C. da R. Neto, C. Luiz, M. J. Rossi and R. M. Di Piero (2017) *Lactobacillus plantarum* exopolysaccharides induce resistance against tomato bacterial spot. *Journal of Agricultural Science* 9:162–179, <https://doi.org/10.5539/jas.v9n2p162>
- Blainski J. M. L., A. C. da R. Neto, E. C. Schmidt, J. A. Voltolini, M. J. Rossi and R. M. Di Piero (2018) Exopolysaccharides from *Lactobacillus plantarum* induce biochemical and physiological alterations in tomato plant against bacterial spot. *Applied Microbiology and Biotechnology* 102:4741–4753, <https://doi.org/10.1007/s00253-018-8946-0>
- Boller T. and G. Felix (2009) A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern-recognition receptors. *Annual Review of Plant Biology* 60:379–406, <https://doi.org/10.1146/annurev>

- arplant.57.032905.105346
- Bradford M. M. (1976) A rapid and sensitive method for the quantitation microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry* 72:248–254, [https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3)
- Cavalcanti F. R., M. L. V. Resende, C. P. S. Carvalho, J. A. G. Silveira and J. T. A. Oliveira (2006) Induced defense responses and protective effects on tomato against *Xanthomonas vesicatoria* by an aqueous extract from *Solanum lycocarpum* infected with *Crinipellis perniciosa*. *Biological Control* 39:408–417, <https://doi.org/10.1016/j.biocontrol.2006.05.009>
- Cavalcanti F. R., M. L. V. Resende, C. P. S. Carvalho, J. A. G. Silveira and J. T. A. Oliveira (2007) An aqueous suspension of *Crinipellis perniciosa* mycelium activates tomato defense responses against *Xanthomonas vesicatoria*. *Crop Protection* 26:729–738, <https://doi.org/10.1016/j.cropro.2006.06.012>
- Dewen Q., D. Yijie, Z. Yi, L. Shupeng and S. Fachao (2017) Plant immunity inducer development and application. *Molecular Plant-Microbe Interactions* 30:355–360, <https://doi.org/10.1094/mpmi-11-16-0231-cr>
- Gottig N., C. V. Vranich, G. G. Sgro, A. Piazza and J. Ottado (2018) HrpE, the major component of the *Xanthomonas* type three protein secretion pilus, elicits plant immunity responses. *Scientific Reports* 8:1–13, <https://doi.org/10.1038/s41598-018-27869-1>
- Huang C. H. and G. E. Vallad (2018) Soil applications of acibenzolar-S-methyl induce defense gene expression in tomato plants against bacterial spot. *European Journal of Plant Pathology* 150:971–981, <https://doi.org/10.1007/s10658-017-1336-0>
- Jones J. B., G. H. Lacy, H. Bouzar, R. E. Stall and N. W. Schaad (2004) Reclassification of the *Xanthomonas* associated with bacterial spot disease of tomato and pepper. *Systematic and Applied Microbiology* 27:755–762, <https://doi.org/10.1016/j.syapm.2005.05.002>
- Laemmli U. K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 228:680–685.
- Lanna-Filho R., R. M. Souza, M. M. Magalhães, L. Villela, E. Zanotto, P. M. Ribeiro-Júnior and M. L. V. Resende (2013) Induced defense responses in tomato against bacterial spot by proteins synthesized by endophytic bacteria. *Tropical Plant Pathology* 38:295–302, <https://doi.org/10.1590/S1982-56762013005000011>
- Louws F. J., M. Wilson, H. L. Campbell, D. A. Cuppels, J. B. Jones, P. B. Shoemaker, F. Sahin and S. A. Miller (2001) Field control of bacterial spot and bacterial speck of tomato using a plant activator. *Plant Disease* 85:481–488, <https://doi.org/10.1094/pdis.2001.85.5.481>
- Luiz C., A. C. Rocha-Neto and R. M. Di-Piero (2015) Resistance to *Xanthomonas gardneri* in tomato leaves induced by polysaccharides from plant or microbial origin. *Journal of Plant Pathology* 97:119–127, <https://doi.org/10.4454/JPP.V97I1.029>
- Mishina T. E. and J. Zeier (2007) Pathogen-associated molecular pattern recognition rather than development of tissue necrosis contributes to bacterial induction of systemic acquired resistance in *Arabidopsis*. *Plant Journal* 50:500–513, <https://doi.org/10.1111/j.1365-3113X.2007.03067.x>
- Murashige T. and F. Skoog (1962) A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiologia Plantarum* 15:473–497, <https://doi.org/10.1111/j.1399-3054.1962.tb08052.x>
- Potnis N., S. Timilsina, A. Strayer, D. Shantharaj, J. D. Barak, M. L. Paret, G. E. Vallad and J. B. Jones (2015) Bacterial spot of tomato and pepper: diverse *Xanthomonas* species with a wide variety of virulence factors posing a worldwide challenge. *Molecular Plant Pathology* 16:907–920, <https://doi.org/10.1111/mpp.12244>
- Ritchie D. F. (2000) Bacterial spot of pepper and tomato. The plant health instructor, <https://www.apsnet.org/edcenter/disandpath/prokaryote/pdlessons/Pages/Bacteriaspot.aspx>
- Reimer-Michalski E. M. and U. Conrath (2016) Innate immune memory in plants. *Seminars in Immunology* 28:319–327, <https://doi.org/10.1016/j.smim.2016.05.006>
- SIAP, Servicio de Información Agroalimentaria y Pesquera (2021). Anuario Estadístico de la Producción Agrícola. Retrieved from Servicio de Información Agroalimentaria y Pesquera, <https://nube.siap.gob.mx/cierreagricola/>
- Šević M., K. Gašić, M. Ignjatov, M. Mijatović, A. Prokić and A. Obradović (2019) Integration of biological and conventional treatments in control of pepper bacterial spot. *Crop Protection* 119:46–51, <https://doi.org/10.1016/j.cropro.2019.01.006>
- Sharma S. and K. Bhattarai (2019) Progress in developing bacterial spot resistance in tomato. *Agronomy* 9:1–11, <https://doi.org/10.3390/agronomy9010026>
- Strayer A. L., A. Jeyaprakash, G. V. Minsavage, S. Timilsina, G. E. Vallad, J. B. Jones and M. L. Paret (2016) A multiplex real-time PCR assay differentiates four *Xanthomonas* species associated with bacterial spot of tomato. *Plant Disease* 100:1660–1668, <https://doi.org/10.1094/pdis-09-15-1085-re>
- Thayer P. L. and R. E. Stall (1961) A survey of *Xanthomonas vesicatoria* resistance to streptomycin. *Proceedings of the Florida State Horticultural Society* 75:163–165.
- Vallad G. E., K. L. Pernezny, B. Balogh, A. Wen, J. F. L. Figueiredo, J. B. Jones, T. Momol, R. M. Muchovej, N. Havranek, N. Abdallah, S. Olson and P. D. Roberts (2010) Comparison of kasugamycin to traditional bactericides for the management of bacterial spot on tomato. *HortScience* 45:1834–1840, <https://doi.org/10.21273/HORTSCI.45.12.1834>
- Vallad G., K. Pernezny and T. Momol (2004) A series on diseases in the Florida vegetable Garden: tomato. Retrieved from University of Florida, IFAS Extension, <https://edis.ifas.ufl.edu/pdf/PP/PP12100.pdf>
- Voets L., H. D. Boulois, L. Renard, D. G. Strullu and S. Declerck (2005) Development of an autotrophic culture system for the in vitro mycorrhization of potato plantlets. *FEMS Microbiology Letters* 248:111–118, <https://doi.org/10.1016/j.femsle.2005.05.025>
- Wu S., L. Shan and P. He (2014) Microbial signature-triggered plant defense responses and early signaling mechanisms. *Plant Science* 228:118–126, <https://doi.org/10.1016/j.plantsci.2014.03.001>
- Zipfel C. (2014) Plant pattern-recognition receptors. *Trends in Immunology* 35:345–351, <https://doi.org/10.1016/j.it.2014.05.004>