



ALLOMETRIC EQUATIONS FOR BELOWGROUND BIOMASS AND CARBON CONTENT OF *Pinus patula* SCHL. et CHAM.

ECUACIONES ALOMÉTRICAS PARA BIOMASA Y CONTENIDO DE CARBONO SUBTERRÁNEO DE *Pinus patula* SCHL. et CHAM.

Frinet Medrano-Meraz^{1*}, Miguel Á. López-López¹, Gregorio Ángeles-Pérez¹, Francisco Cruz-Cobos² y Bernardus H. Jozeph-de-Jong³

¹Colegio de Postgraduados, Postgrado en Ciencias Forestales, Montecillo, Texcoco, Edo. de México, México. ²Instituto Tecnológico de El Salto, División de Estudios de Posgrado e Investigación, El Salto, Durango, México. ³El Colegio de la Frontera Sur, Departamento de Ciencias de la Sustentabilidad, Lerma, Campeche, México.

*Autor de correspondencia (medrano.frinet@colpos.mx)

SUMMARY

Allometric equations are essential tools for estimating biomass and carbon fixation in forest ecosystems. Published information is limited for belowground biomass of forests, mainly due to the methodological challenges of its determination. With the aim of contributing to the study of this important carbon store, in the present study allometric equations were developed for the estimation of belowground biomass of *Pinus patula* Schiede ex Schldl. et Cham. in its main natural distribution area: the states of Hidalgo, Puebla and Veracruz, Mexico. Seven *Pinus patula* trees with varying dimensions were selected and torn down at various sites; their diameter at breast height (DBH), total height (H), and crown diameter (CD) were measured. The totality of fine and coarse roots was extracted; the aboveground biomass was quantified by census, by sampling or by means of an allometric equation. Equations were developed for belowground biomass and carbon content, both total and by components, as well as for individual root, both total and by components. The average root/shoot ratio was 0.20 and the proportion of belowground in relation to the total biomass ranged from 12.1 to 21.2 %. The data on belowground biomass and carbon content at the tree level better fitted the Schumacher and Hall model that included both DBH and H as independent variables, which explained 99 % of the variation of belowground biomass and carbon. It was concluded that it is possible to use allometric equations to estimate the belowground biomass and carbon of *Pinus patula* trees from DBH and H, or even using only DBH.

Index words: *Pinus patula*, aboveground biomass, carbon fixation, coarse roots, fine roots, root/shoot ratio.

RESUMEN

Las ecuaciones alométricas son herramientas esenciales para estimar la biomasa y la fijación de carbono en los ecosistemas forestales. La información publicada es limitada para la biomasa subterránea de bosques, debido a los desafíos metodológicos que representa su determinación. Con el objetivo de contribuir al estudio de este importante almacén de carbono, en el presente estudio se desarrollaron ecuaciones alométricas para la estimación de biomasa subterránea de *Pinus patula* Schiede ex Schldl. et Cham. en el área principal de distribución natural: los estados de Hidalgo, Puebla y Veracruz, México. Se seleccionaron y derribaron en diversos sitios siete árboles de *Pinus patula* con dimensiones variadas; se midió su diámetro a la altura de pecho (DAP), altura total (AT) y diámetro de copa (DC). Se extrajo la totalidad de las raíces finas y gruesas; la biomasa aérea se cuantificó por censo, muestreo o mediante una ecuación alométrica. Se desarrollaron ecuaciones

para biomasa subterránea y contenido de carbono totales y por componentes, así como para raíz individual total y por componentes. El cociente de raíz/parte aérea promedio fue de 0.20 y la proporción de biomasa subterránea con respecto a la biomasa total por árbol varió de 12.1 a 21.2 %. Los datos de biomasa subterránea y contenido de carbono total a nivel de árbol se ajustaron mejor al modelo Schumacher y Hall, que incluye el DAP y AT como variables independientes, y explicaron el 99 % de la variación en biomasa y carbono subterráneo. Se concluyó que es posible usar ecuaciones alométricas para estimar la biomasa y carbono subterráneo de árboles de *Pinus patula* a partir del DAP y AT, o incluso usando solo el DAP.

Palabras clave: *Pinus patula*, biomasa aérea, cociente raíz/parte aérea, fijación de carbono, raíces finas, raíces gruesas.

INTRODUCTION

Allometric equations are essential tools for estimating forest biomass and carbon stocks in forests (Tashi *et al.*, 2017). These estimates make it possible to evaluate the impact of forests on climate change mitigation (Ngoma *et al.*, 2018) and, henceforth, to support the adoption of environmental mechanisms and policies (Agrawal *et al.*, 2011). These type of equations generally express the relationships between tree biomass and the diameter at breast height (DBH), tree height or any other easy-to-measure dendrometric variables in forest inventories (Fonseca *et al.*, 2009; Henry *et al.*, 2013; Ketterings *et al.*, 2001; Picard *et al.*, 2015; Razakamanarivo *et al.*, 2012).

Ideally, *ad hoc* allometric equations should be used for local conditions and species in order to minimize prediction errors (Watzlawick *et al.*, 2001; Zapata-Cuarteros *et al.*, 2012). According to Cole and Ewel (2006), application of generic equations to specific sites or species frequently results in uncertain estimations. Analogously, Chave *et al.* (2014) and Koala *et al.* (2017) agree that the use of species-specific equations gives rise to lower uncertainties.

Globally, multiple equations for aboveground biomass

prediction have been developed by well-established methods; whereas, few belowground biomass equations have been established (Cairns *et al.*, 1997; Razakamanarivo *et al.*, 2012). The belowground component of trees is the least studied one, due to the methodological difficulties, that result in slow and expensive processes (Brown, 1997; Návar, 2015). Moreover, there is no agreement on accurate methods to be used for such a challenge (Addo-Danso *et al.*, 2016; Brown, 2002; Gárate and Blanco, 2013).

Nevertheless, belowground biomass is an important carbon pool that should be included in the local, regional and global carbon estimations (Varik *et al.*, 2013), since it may represent up to 40% of green biomass in conifer forests (Litton *et al.*, 2003). Accordingly, in order to contribute information on belowground biomass, this study aimed to develop allometric equations for estimating belowground biomass and carbon content, both at the whole-tree level and at the individual-root level, for *Pinus patula*, which is an important tree species from the ecological and economic viewpoints.

MATERIALS AND METHODS

Study area

The present study was carried out at the municipalities of Huayacocotla, Veracruz; Acaxochitlán, Hidalgo; Zacualtipan, Hidalgo; Chignahuapan, Puebla; and Zacatlan, Puebla, Mexico, between the coordinates 19° 46' 05"-20° 37' 8.7" N and 98° 00' 21.9"-98° 36' 29.3" W. The mean elevation of the sampled sites varied from 2055 to 2600 masl. *Pinus patula* Schiede ex Schldl. et Cham. is a species native to Mexico, it grows from 17 to 24° latitude N (Perry, 1991; Vela, 1980). Hidalgo, Puebla and Veracruz are the states where the most abundant populations are found. According to Vela (1980), the lowest temperatures along the *Pinus patula* natural distribution area occur during the months of December or January. The highest temperatures take place in May. Minimum temperatures vary from 0.9 to 7.9 °C and the highest ones fluctuate from 20.3 to 27.2 °C. Sites where *P. patula* grows are moist and have good drainage, with annual precipitation varying from 1000 to 1500 mm. *P. patula* grows in pure stands or it may associate with *P. greggii* Engelm., *P. maximinoi* H. E. Moore, *P. pseudostrobus* Lindl., *P. teocote* Schiede. ex Schldl. et Cham., *P. leiophylla* Schiede ex Schldl. et Cham., *P. patula* Schiede ex Schldl. et Cham. var. *longipedunculata* Looock ex Martinez, *Abies religiosa* (Kunth) Schldl. et Cham., *P. montezumae* Lamb., *P. ayacahuite* Ehrenb. ex Schldl. var. *veitchii* (Roetzl) Shaw, and *Quercus* spp. (Perry, 1991). Soils vary from moderately to very deep, with loamy texture in the shallow horizons and clayey texture in the deepest horizons (Vela, 1980).

Sampling procedure

The study aimed to develop two groups of allometric equations: 1) tree-level belowground biomass equations, and 2) individual-root level biomass equations. In both cases, a destructive sampling of trees was used, which consisted of extracting the roots from selected healthy trees, with a single straight stem. Trees were dominant in their populations and were strategically located throughout the species natural distribution area, in order to assure their representativeness. Root extraction was conducted from June 2016 to September 2017.

For the first group of equations, seven *P. patula* trees were selected in the following sites: two trees at Huayacocotla, Veracruz; two trees at Chignahuapan, Puebla; one tree at Zacatlán, Puebla; one tree at Acaxochitlán, Hidalgo and one tree at Zacualtipan, Hidalgo. The DBH ranged from 1.5 to 57.0 cm. Due to the high degree of difficulty and the high costs involved in root extraction, it was only carried out in this number of trees, which is intended to represent the entire main natural distribution area of this species, and also the most common range of tree dimensions. There are no reports on belowground biomass equations for large *P. patula* trees, that is why large trees were included, despite selecting a small sample size. In other studies, successful belowground allometric equations for small trees have been developed by using sample sizes similar or even lower than the one used in this study (Xiao *et al.*, 2003; Xiao and Ceulemans, 2004), or by using two or three trees per diameter class (Rathore *et al.*, 2018).

To develop the individual-root set group of equations, 11 trees were selected, from which 66 roots with basal diameter (RBD) ranging from 6.82 to 155.0 cm were extracted. Some of the roots came from the aforementioned seven trees, but many other roots were extracted from independent trees also located in the mentioned localities, in order to assure a wide morphological variability of the roots.

Dasometric variables measured

Variables measured on each of the standing trees were: 1) DBH in cm; this variable was measured on standing trees at a height of 1.3 m above ground level by using a diameter tape; 2) crown diameter (CD) in m, this variable was measured by using a metric tape; 3) after felling off the tree, its total height in m was determined with a metric tape. As for the individual roots, the RBD as well as the total root length (RL; cm) were measured in the field, after extracting the whole root; for large roots (RBD > 5 cm) the RBD was measured by using a diameter tape; for small roots (RBD ≤ 5 cm) a caliper was used.

Belowground biomass and carbon content

To determine the root biomass, each of the seven selected trees were felled; then, with conventional agricultural and gardening tools (shovels, picks, hoes, trowels, etc.), the coarse roots around the tree trunk were identified and marked. Thereafter, excavation along each individual root was continued, until exposing the whole root, including both the coarse and the fine components. In this process, when a root portion, whatever the order it was, got accidentally excised, it was recollected, and the excision point labeled for later integration. By using this procedure, the totality of the tree roots, including the tap root was extracted. Regarding individual-root biomass, from each of the 11 selected trees one or several roots were exposed and extracted by following the procedure explained before.

In laboratory, roots were cleaned and classified as primary, secondary and tertiary or higher order roots (Atkinson, 2000); moreover, each of these groups were reclassified into coarse (root diameter > 2 mm) or fine roots (root diameter ≤ 2 mm) (IPCC, 2006). Each group was placed into perfectly labeled paper bags that were introduced into a forced-air oven (Sheldon Model 1675-S®, Cornelius, Oregon, USA) calibrated to 70 °C. Sample bags containing the biggest pieces of root wood were weighted daily and all the bags were taken out the oven when weight of sample materials became constant. Thereafter, the bags were weighted by using an OHAUS scale 2 kg in capacity and 1 g in precision (OHAUS Corp., Parsippany, New Jersey, USA). The classification process gave rise to the following root components: 1) tap root (TaR), is the root originating from the tree root collar, generally goes down into deep soil horizons; this root gives rise to the lateral roots; 2) primary roots (PR), come directly from the tap root; 3) secondary roots (SR), emerging from the primary roots; 4) tertiary roots (TR), emerge from the secondary roots; 5) coarse roots (CR), include all-order roots with diameter ≥ 2 mm; 6) fine roots (FR), include all-order roots with diameter < 2 mm; 7) coarse primary roots (CPR); 8) fine primary roots (FPR); 9) coarse secondary roots (CSR); 10) fine secondary roots (FSR); 11) coarse tertiary roots (CTR), and 12) fine tertiary roots (FTR).

The belowground carbon content of the seven-sample trees was determined by multiplying the total belowground biomass of each of the trees by the average carbon concentration for *Pinus patula* reported by Diaz-Franco *et al.* (2007).

Aboveground biomass

Aboveground biomass, that is, the biomass of stems,

branches and foliage of the seven trees processed was quantified by using the method that, according to the size of the trees, would optimize time and cost, as well as precision of the estimates. In the case of small trees (DBH < 10 cm), after splitting the aboveground components, all the biomass was processed (dried) in laboratory. Aboveground biomass of the 16.4 cm DBH tree was determined by sampling using the dry weight/fresh weight ratio. First, the fresh weight of each of the structural components was obtained with an OHAUS® AS-100 scale 100 kg in capacity and 100 g in precision (OHAUS Corp., Parsippany, New Jersey, USA). The fresh weight of 9 to 12 random samples per component was also recorded. The samples were subsequently dried in laboratory in a Sheldon Model 1675-S® forced air-circulation oven at 70 °C until constant weight. Once both weights were obtained, the moisture content of the sampling units was determined (Equation 1; Flores-Nieves *et al.*, 2011) and then, the dry weight of components was computed by applying Equation 2. Finally, the total aboveground biomass of the tree was estimated by integrating the biomass of the several structural components (Domínguez-Cabrera *et al.*, 2009). The aboveground biomass of the structural components of trees with DBH higher than 25.5 cm was estimated by the allometric equations reported by Castellanos *et al.* (1996) for *P. patula*.

$$MCs = \frac{(fws - dws) \cdot 100}{fws} \quad (1)$$

$$DWc = \frac{FWc \cdot (100 - MCm)}{100} \quad (2)$$

where *MCs* is the moisture content of the sampling unit (%), *fws* is the fresh weight of the sampling unit (kg), *dws* is the dry weight of the sampling unit (kg), *DWc* is the dry weight of the tree component (kg), *FWc* is the fresh weight of the tree component (kg) and *MCm* is the mean moisture content of the component sample (%).

After obtaining the total below and aboveground biomass in the seven harvested trees, the root/shoot ratio was determined, as well as the proportion of the below and aboveground biomass in relation to the total biomass. Table 1 summarizes the variables measured for each observation.

The RBD for the 66 individual roots ranged from 6.8 to 155.0 mm and length varied from 60 to 2173 cm. Total root biomass (ToRB) displayed the highest variation (*s* = 3967.8 g), followed by CR (*s* = 3956.9 g). Only five out of 66 roots extracted had fine primary roots, indicating that this type of roots is infrequent in *P. patula* trees (Table 2).

Statistical analysis

The data obtained from belowground biomass, belowground carbon content, and the dasometric variables measured on the seven sampled trees, allowed to test models that are frequently used for tree allometry. The independent variables H, CD and DBH were tested one by one. The DBH showed the highest association with the dependent variable (belowground biomass). In the scatter plot of the observed values of belowground biomass for *P. patula* (Figure 1) with respect to DBH, a potential relationship was determined. Fonseca *et al.* (2009) suggested that DBH

is the variable most strongly associated with biomass.

The models that more appropriately described the dispersion of data were selected, trying those models that were simple and more frequently cited in the relevant literature on the subject (Table 3). Based on those models, allometric equations were developed for the prediction of total and by- component belowground biomass and the corresponding carbon content at the tree level, as a function of DBH and H or simply DBH as the independent variable. On the other hand, biomass data for individual roots fitted better to model 3 (Table 3). The allometric equations to

Table 1. Summary of the variables measured in the *Pinus patula* sampled trees.

Tree	DBH (cm)	H (m)	CD (m)	Belowground biomass (kg)	Aboveground biomass (kg)	Total biomass (kg)	Belowground carbon content (kg)	R	B/T (%)	A/T (%)
1	1.5	2.5	1.2	0.05	0.39	0.44	0.03	0.13	12.1	88.0
2	9.0	9.8	2.7	3.19	16.20	19.38	1.60	0.20	16.4	83.6
3	16.4	21.8	2.5	13.13	85.23	98.37	6.61	0.15	13.4	86.7
4	25.5	16.6	6.4	61.12	247.70	308.81	30.75	0.25	19.8	80.2
5	35.5	20.7	9.3	111.39	528.33	639.72	56.04	0.21	17.4	82.6
6	46.7	20.6	--	196.41	990.43	1186.84	98.81	0.20	16.6	83.5
7	57.0	32.5	8.0	420.66	1564.29	1984.95	211.63	0.27	21.2	78.8
Mean	27.4	17.8	4.3	115.14	490.37	605.50	57.92	0.20	16.7	83.3

DBH: diameter at breast height, H: total height, CD: crown diameter, R: root/shoot ratio, B/T: belowground biomass/total biomass ratio, A/T: aboveground biomass/total biomass ratio.

Table 2. Descriptive statistics of individual roots by components.

Variable	N	Minimum	Mean	Maximum	SD
RBD (mm)	66	6.82	62.70	155	38.20
RL (cm)	64	60.00	469.49	2173	403.04
ToR (g)	66	12.16	3194.62	18152	3967.83
PR (g)	60	7.30	2255.63	14769	2786.67
SR (g)	59	3.00	931.23	8116	1458.14
TR (g)	58	0.20	248.96	2468	453.83
CR (g)	66	10.31	3173.84	18090	3956.92
FR (g)	65	0.94	21.10	116	20.86
CPR (g)	60	7.30	2255.47	14769	2786.78
FPR (g)	5	0.33	1.87	3	1.19
CSR (g)	59	2.00	926.85	8113	1458.53
FSR (g)	49	1.00	5.28	19	4.40
CTR (g)	48	0.16	279.57	2411	469.21
FTR (g)	58	0.20	17.59	110	19.12343

N: number of roots, SD: standard deviation, RBD: root basal diameter, RL: total root length, ToR: total root biomass, PR: primary roots, SR: secondary roots, TR: tertiary roots, CR: coarse roots, FR: fine roots, CPR: coarse primary roots, FPR: fine primary roots, CSR: coarse secondary roots, FSR: fine secondary roots, CTR: coarse tertiary roots, FTR: fine tertiary roots.




Figure 1. Dispersion of values and trend line for belowground biomass in relation to diameter at breast height (DBH).

estimate total and by-component biomass of individual roots, used only RBD as the independent variable. Also, as a function of this variable, a special equation for the tap root and its components was developed, since such tree structure exhibits a particular morphometry.

Model fitting was carried out by linear and nonlinear regression using the MODEL and REG procedures of the Statistical Analysis System (SAS) version 9.4 for Windows (SAS Institute, 2011). The parameters used to evaluate model goodness of fit was the adjusted R^2 , which provides the variability of the dependent variables explained by the model, considering the number of parameters, the root mean square error (RMSE) and the Akaike's Information Criterion (AIC), which is frequently used for comparisons between models (Akaike, 1974; Picard *et al.*, 2012). The relative error was calculated using equation 3 (Djomo *et al.*, 2010). In the case of individual root models, the goodness of fit was evaluated based on the coefficient of

determination (R^2) and RMSE.

$$RE = \frac{Pi - Mi}{Mi} \cdot 100 \tag{3}$$

where RE is the relative error (%), Pi is the predicted biomass for the i^{th} tree (kg), Mi is the observed biomass for the i^{th} tree (kg).

RESULTS AND DISCUSSION

Tree-level belowground biomass and carbon content equations

One- and two-input models were used in this study; that is, the regression equations produced for belowground and carbon content biomass estimation tree-level used DBH and H as independent variables, in order to take advantage of the contribution of H. In both cases, according to the evaluation statistics (RMSE, AIC, RE and adjusted R^2), model 1, which included H as an additional independent variable, resulted in lower values for RMSE, AIC and RE, and higher adjusted R^2 (0.99). In contrast, the adjusted R^2 for model 2 was 0.98 (Table 4; Table 5). Model 1 tended to overestimate belowground biomass (RE = 6.1 %), while model 2 underestimated it (RE = -28 %; Table 4). Similar results were obtained for carbon content, with mean relative errors of 4.9 and -32 %, respectively (Table 5). The mean relative error is considerably lower in both estimations when model 1, which included DBH and H as independent variables, was used. Djomo *et al.* (2010) reported contrasting results for aboveground biomass estimations since they got a mean relative error

Table 3. Models selected to estimate belowground biomass and carbon content of *Pinus patula*.

Model	Equation	Author
1	$y = \beta_0 DBH^{\beta_1} H^{\beta_2}$	Schumacher and Hall (1933)
2	$y = \beta_0 DBH^{\beta_1}$	Ter-Mikaelian and Korzukhin (1997)
3	$\ln(y) = \ln(\beta_0) + \beta_1 \ln(RBD)$	Baskerville (1972)

y : belowground biomass or belowground biomass component (kg), DBH : diameter at breast height (cm), H : tree total height (m), RBD : root basal diameter (mm) β_0 , β_1 , and β_2 : parameters to be estimated, \ln : natural logarithm.

of 7.4 % when used DBH only as the independent variable; however, when they added tree height or wood density as independent variables, the estimations did not improve significantly.

Nam *et al.* (2016) pointed out that allometric models should not only be precise, but also as simple as possible. Besides of being an easy to measure variable, DBH is preferred as an independent variable because of the lower uncertainty it brings about as compared to tree height, which is a hard-to-measure variable (López, 2005). Peichl and Arain (2007) suggested that DBH as an independent variable is sufficient to precisely estimate above- and belowground biomass; however, it is important to consider the error of estimations. In this study, even though model 2 (which included only DBH as the independent variable) showed an appropriate fitting to data, model 1 (whose independent variables are DBH and H) resulted in significantly lower relative error.

When the belowground biomass, estimated with models 1 and 2, was plotted against the observed belowground biomass, the values looked quite close to the regression line, which suggests a good fit in both cases. Similarly, slopes and determination coefficients for the linear regression were nearly 1 ($R^2 = 0.99$ for model 1 and $R^2 = 0.98$ for model 2). According to Figure 2, model 1 makes better predictions of total biomass than model 2.

Tree-level belowground biomass equations by components

In general, biomass of the most abundant root components, CPR and FTR, is more closely linked to DBH, while that of the least frequent components, FPR and FSR, exhibits the lowest dependence to DBH. For model 2, R^2 values were 0.95 and 0.86 for CTR and FTR, respectively, while for the same components the adjustment parameters of model 1 were significant with higher determination coefficients that reached 0.99 (Table 6). Correlations between FR and DBH or H were too low, probably because quantification of FR included biomass of the fine primary roots, a variable that showed extremely small values, regardless of tree size. Perhaps an increase in the sample size (addition of sampled trees) could allow an improved model fitting. It was not possible to generate models for FR and FPR (Table 5). When fitting preliminary models, Mugasha *et al.* (2013) and Liepins *et al.* (2018) recommend to try little improvements in model fitting. This seems to be the case of this study, since it shows that model 1 when adjustment was possible, resulted in R^2 values higher than 0.90. The use of DBH alone as the independent variable is likely to be sufficient to explain the variation of the root biomass components, except for FSR and components including tertiary roots; in those cases, only the use of both independent variables allowed to parametrize the models or to improve model fitting.

Table 4. Model parameters and statistics of goodness of fit to estimate total belowground biomass for *Pinus patula*.

Model	Estimated parameters	SE	Pr > t	adj-R ²	AIC	RMSE (kg)	RE (%)
1	β_0	0.0074	0.0035	0.0979	0.99	33.87	8.40
	β_1	2.0780	0.2061	0.0005			
	β_2	0.7294	0.1623	0.0109			
2	β_0	0.0023	0.0027	0.4212	0.98	44.53	18.55
	β_1	2.9883	0.2872	0.0001			

SE: standard error, adj-R²: adjusted coefficient of determination, AIC: Akaike's information criterion, RMSE: root mean square error, RE: relative error.

Table 5. Model parameters and statistics of goodness of fit to estimate belowground carbon content for *Pinus patula*.

Model	Estimated parameters	SE	Pr > t	adj-R ²	AIC	RMSE (kg)	RE (%)
1	β_0	0.0037	0.0017	0.0979	0.99	24.25	4.22
	β_1	2.0780	0.2061	0.0005			
	β_2	0.7294	0.1623	0.0109			
2	β_0	0.0011	0.0013	0.4214	0.98	34.91	9.33
	β_1	2.9888	0.2873	0.0001			

SE: standard error, adj-R²: adjusted coefficient of determination, AIC: Akaike's Information Criterion, RMSE: root mean square error, RE: relative error.




Figure 2. Relationship between observed and predicted belowground biomass. A) Estimated values by model 1, B) Estimated values by model 2.

Biomass equations by components for individual roots

Several equations for individual root components were derived from data collected from 66 individual roots. The equations presenting the best fitting were ToR, PR, SR, CR, CPR and CSR; all of them showed R^2 values higher than 0.7. All the equation parameters were significant ($P \leq 0.05$), except those for FR, FPR, and FSR which, consequently, experienced the lowest determination coefficients (Table 7).

More than 90 % of the observed variability in ToR, PR, TR, CR, CPR, FSR and CTR was explained by the basal diameter of the tap root. In the case of CSR, correlation with tap-root basal diameter also was significant, although tap-root basal diameter explained a lower proportion of its variability. The equations that had non-significant parameters were SR, FR and FTR. As fine roots, and particularly fine primary roots along the tap root are scarce, it was not possible to fit the FPR data to a model (Table 8).

In general, better fittings were observed to the belowground biomass data at the tree-level than to the individual-root data. It appears that belowground biomass is highly correlated with DBH; contrarily, dependence of individual-root biomass to its basal diameter is weak, although such correlation could be species-dependent. Ríos-Carrasco and Nívar-Chaidez (2010) found a R^2 of 0.84 for pinion pine root biomass as a function of root basal diameter. Variation of biomass of individual roots of *P. patula* is likely to be explained by factors such as water and nutrient availability, soil compaction or texture, among others.

Below- and aboveground biomass

The root/shoot biomass ratio varied from 0.13 to 0.27, with a mean value of 0.20 (Table 1). In the study conducted by Sanquetta *et al.* (2011), which included two pine species, they reported a belowground/aboveground biomass ratio of 0.17, which is slightly lower than the value obtained in this study. On the other hand, Liepins *et al.* (2018) found higher values than those of this study: 0.20 to 0.25 for *P. sylvestris* L., 0.21 to 0.30 for *Picea abies* L. Karst, 0.23 to 0.34 for *Betula pendula* Roth and 0.21 to 0.39 for *Populus tremula* L. In 10- and 20-years old *P. ponderosa* Dougl. ex Laws plantations, the belowground to aboveground biomass ratios ranged from 0.21 to 0.24 for the sampled trees; however, for the whole stands, the values varied from 0.23 to 0.25. In both cases, the highest mean values were registered in the younger plantation (Laclau, 2003). This situation is similar to that found by Peichl and Arain (2007) in Canada, who determined that the root/aboveground biomass ratio decreased as the stand age of *P. strobus* L. increased.

In this study, the proportion of belowground biomass in relation to total biomass ranged between 13.4 and 21.2 %, being 16.7 % the mean value (Table 1). This value is lower than that reported for *Vochysia guatemalensis* Donnell-Smithii, where the belowground biomass reached 18.6 % of the total tree biomass (Fonseca *et al.*, 2012). In the study conducted by Martínez *et al.* (2016) the proportion of root biomass in relation to total biomass of *P. cooperi* B. Blanco (eight to 13 years old and 7.6 to 12.2 cm DBH) was 17.57 %, thus, being a value quite similar to the observed in this

Table 6. Model parameters and goodness of fit statistics to estimate the belowground biomass per component at the tree level.

Component	Model	Estimated parameters	SE	Pr > t	adj-R ²	RMSE (kg)	
TaR (kg)	2	β_0	0.00018	0.0002	0.4369	0.98	6.48
		β_1	3.38595	0.2970	< 0.0001		
PR (kg)	2	β_0	0.00099	0.0027	0.7308	0.91	18.84
		β_1	2.98429	0.6913	0.0076		
SR (kg)	2	β_0	0.00774	0.0066	0.2925	0.98	2.91
		β_1	2.21286	0.2158	0.0002		
TR (kg)	2	β_0	0.00017	0.0003	0.6309	0.95	1.36
		β_1	2.83178	0.4927	0.0022		
CR (kg)	2	β_0	0.00374	0.0070	0.6156	0.95	20.33
		β_1	2.74436	0.4712	0.0021		
CPR (kg)	2	β_0	0.00099	0.0027	0.7309	0.91	18.84
		β_1	2.98448	0.6915	0.0076		
CSR (kg)	2	β_0	0.00685	0.0062	0.3171	0.98	3.03
		β_1	2.24293	0.2284	0.0002		
CTR (kg)	2	β_0	0.00009	0.0002	0.6422	0.95	1.24
		β_1	2.97466	0.5092	0.0021		
FTR (kg)	2	β_0	0.00130	0.0025	0.6232	0.86	0.17
		β_1	1.66772	0.4906	0.0193		
TR (kg)	1	β_0	0.0000043	0.0000031	0.2477	0.99	0.25
		β_1	5.10711	0.2999	< 0.0001		
		β_2	-1.59354	0.1549	0.0005		
CTR (kg)	1	β_0	0.0000012	0.0000011	0.3368	0.99	0.25
		β_1	5.46879	0.3705	0.0001		
		β_2	-1.67034	0.1855	0.0008		
FSR (kg)	1	β_0	0.00085	0.0015	0.6014	0.96	0.06
		β_1	-2.95403	0.4221	0.0022		
		β_2	4.92840	0.8096	0.0037		
FTR (kg)	1	β_0	0.00094	0.0004	0.1242	0.99	0.03
		β_1	3.21843	0.2113	0.0001		
		β_2	-1.75394	0.1595	0.0004		

SE: standard error, adj-R²: adjusted coefficient of determination, RMSE: root mean square error, TaR: tap root, PR: primary roots, SR: secondary roots, TR: tertiary roots, CR: coarse roots, CPR: coarse primary roots, CSR: coarse secondary roots, FSR: fine secondary roots, CTR: coarse tertiary roots, FTR: fine tertiary roots.

study, even though in that case, dimensions of the trees were lower than those in this research.

The proportion of belowground biomass in relation to total biomass in *P. patula* trees increased as the DBH went up (Figure 3). This trend coincides with that reported by Grier *et al.* (1981) for *Abies amabilis* Douglas ex J. Forbes stands, and also with that of Magnani *et al.* (2000), who registered an increased biomass of fine roots as the age of

the *P. sylvestris* stands increased from 7 to 59 years.

Nevertheless, another study on *P. sylvestris* reported a decrease in this proportion as the trees got larger (Kenina *et al.*, 2018), which means that the B/T ratio may be determined not only by the species, but also by the site conditions, since root extraction methodology was similar to the employed in this study, except that fine root biomass was not determined for *P. sylvestris*.

Table 7. Model parameters and goodness of fit statistics to estimate the belowground biomass by component at the individual root level.

Component	Model	Estimated parameters	SE	Pr > t	R ²	RMSE (g)	
ln (ToR; g)	3	β_0	-3.25206	0.4230	< 0.0001	0.90	0.64
		β_1	2.57339	0.1059	< 0.0001		
ln (PR; g)	3	β_0	-2.94077	0.5147	< 0.0001	0.86	0.67
		β_1	2.41009	0.1266	< 0.0001		
ln (SR; g)	3	β_0	-4.39779	0.7805	< 0.0001	0.74	1.02
		β_1	2.46898	0.1922	< 0.0001		
ln (TR; g)	3	β_0	-5.73979	0.9800	< 0.0001	0.64	1.21
		β_1	2.41135	0.2399	< 0.0001		
ln (CR; g)	3	β_0	-3.49708	0.4323	< 0.0001	0.90	0.66
		β_1	2.62623	0.1083	< 0.0001		
ln (FR; g)	3	β_0	-0.68053	0.4860	0.1664	0.43	0.74
		β_1	0.85737	0.1219	< 0.0001		
ln (CPR; g)	3	β_0	-2.96208	0.5158	< 0.0001	0.86	0.67
		β_1	2.41485	0.1269	< 0.0001		
ln (FPR; g)	3	β_0	2.68750	0.8520	0.0511	0.58	0.36
		β_1	-0.46453	0.2264	0.1325		
ln (CSR; g)	3	β_0	-5.14299	0.8185	< 0.0001	0.74	1.07
		β_1	2.63212	0.2015	< 0.0001		
ln (FSR; g)	3	β_0	1.53599	0.5615	0.0088	0.0003	0.71
		β_1	0.01692	0.1398	0.9042		
ln (CTR; g)	3	β_0	-9.18095	1.7626	< 0.0001	0.55	1.41
		β_1	3.15863	0.4144	< 0.0001		
ln (FTR; g)	3	β_0	-1.72452	0.5671	0.0036	0.50	0.70
		β_1	1.04305	0.1388	< 0.0001		

SE: standard error, ToR: total root biomass, PR: primary roots, SR: secondary roots, TR: tertiary roots, CR: coarse roots, FR: fine roots, CPR: coarse primary roots, FPR: fine primary roots, CSR: coarse secondary roots, FSR: fine secondary roots, CTR: coarse tertiary roots, FTR: fine tertiary roots.

The belowground biomass and total biomass ratios apparently are determined by the species, but they are affected by the sampling methods of both, belowground and aboveground biomass (Adame *et al.*, 2017; Beets *et al.*, 2007; Brassard *et al.*, 2011). Because of the use of a variety of methodologies for tree root sampling, published allometric equations show variations in their mathematical structure and varying uncertainty in their estimations (Waring and Powers, 2017; Yuen *et al.*, 2016).

CONCLUSIONS

The relationship between belowground biomass and DBH can be described by a potential model. The model that best predicts belowground biomass and carbon content at the tree-level is the one of Schumacher and

Hall that includes both DBH and H as independent variables. When DBH and H are included in the equations to estimate tree level belowground biomass by component, the improvement in fitting is slight. Tree-level equations for belowground biomass components exhibit a better fitting than individual-root equations. Biomass of the most abundant roots, CPR and FTR, highly depends on DBH; in contrast, biomass of the least frequent roots, FPR and FSR, show a lower dependence from DBH. In the sampled trees of *P. patula* between 1.5 and 57.0 cm in DBH, the average root/shoot ratio is 0.20 and the belowground biomass/total biomass ratio (B/T) ranges between 13.4 and 21.2 %.

ACKNOWLEDGEMENTS

Definition of methodologies and reviews of manuscript

Table 8. Model parameters and goodness of fit statistics to estimate biomass of the tap root and components.

Component	Model	Estimated parameters	SE	Pr > t	R ²	RMSE (g)	
ln (ToR; g)	3	β_0	-0.65185	0.4998	<0.0001	0.98	0.35
		β_1	2.97697	0.1483	<0.0001		
ln (PR; g)	3	β_0	-0.90522	0.5927	0.2014	0.98	0.67
		β_1	2.98988	0.1744	<0.0001		
ln (SR; g)	3	β_0	-3.11284	3.4604	0.4633	0.79	0.92
		β_1	2.69974	0.9718	0.1088		
ln (TR; g)	3	β_0	-13.37030	1.8928	0.0195	0.97	0.50
		β_1	5.23915	0.5316	0.0101		
ln (CR; g)	3	β_0	-0.75880	0.4986	0.2027	0.99	0.3469
		β_1	2.99123	0.1467	<0.0001		
ln (FR; g)	3	β_0	-1.43166	1.2584	0.3379	0.73	0.85
		β_1	1.11657	0.3883	0.0638		
ln (CPR; g)	3	β_0	-0.90522	0.5927	0.2014	0.98	0.41
		β_1	2.98988	0.1744	<0.0001		
ln (CSR; g)	3	β_0	-1.98434	1.7350	0.3165	0.85	1.21
		β_1	2.53033	0.5106	0.0077		
ln (FSR; g)	3	β_0	-2.81838	0.7673	0.0668	0.93	0.20
		β_1	1.12295	0.2155	0.0349		
ln (CTR; g)	3	β_0	-15.08499	2.1697	0.0061	0.96	0.63
		β_1	5.75540	0.5905	0.0023		
ln (FTR; g)	3	β_0	-1.38823	1.3761	0.3874	0.67	0.93
		β_1	1.05075	0.4247	0.0897		

SE: standard error, ToR: total root biomass, PR: primary roots, SR: secondary roots, TR: tertiary roots, CR: coarse roots, FR: fine roots, CPR: coarse primary roots, CSR: coarse secondary roots, FSR: fine secondary roots, CTR: coarse tertiary roots, FTR: fine tertiary roots.




Figure 3. Change in the belowground biomass / total biomass ratio with changing Diameter at Breast Height (DBH).

were carried out by all authors. FMM and MALL conducted the field work, laboratory processing and data analysis. We acknowledge all persons who kindly participated in the field activities. We also acknowledge CONACYT (Mexican Council for Science and Technology) for the scholarship awarded to the first author. Our special gratitude to Program for Sustainable Landscapes (US-AID), which contributed partial funds through the USDA International Programs Office and the Northern Research Station.

BIBLIOGRAPHY

- Adame M. F., S. Cherian, R. Reef and B. Stewart-Koster (2017) Mangrove root biomass and the uncertainty of belowground carbon estimations. *Forest Ecology and Management* 403:52-60, <https://doi.org/10.1016/j.foreco.2017.08.016>
- Addo-Danso S. D., C. E. Prescott and A. R. Smith (2016) Methods for estimating root biomass and production in forest and woodland ecosystem carbon studies: a review. *Forest Ecology and Management* 359:332-351, <https://doi.org/10.1016/j.foreco.2015.08.015>
- Agrawal A., D. Nepstad and A. Chhatre (2011) Reducing emissions from deforestation and forest degradation. *Annual Review of Environment and Resources* 36:373-396, <https://doi.org/10.1146/annurev-environ-042009-094508>
- Akaike H. (1974) A new look at the statistical model identification. *IEEE Transactions on Automatic Control* 19:716-723, <https://doi.org/10.1109/TAC.1974.1100705>
- Atkinson D. (2000) Root characteristics: why and what to measure. In: *Root Methods*. A. L. Smit, A.G. Bengough, C. Engels, M. van Noordwijk, S. Pellerin and S. C. van de Geijn (eds). Springer. Heidelberg, Germany. pp:1-32, https://doi.org/10.1007/978-3-662-04188-8_1
- Baskerville G. L. (1972) Use of logarithmic regression in the estimation of plant biomass. *Canadian Journal of Forest Research* 2:49-53, <https://doi.org/10.1139/x72-009>
- Beets P. N., S. H. Pearce, G. R. Oliver and P. W. Clinton (2007) Root/shoot ratios for deriving below-ground biomass of *Pinus radiata* stands. *New Zealand Journal of Forestry Science* 37:267-288.
- Brassard B. W., H. Y. Chen, Y. Bergeron and D. Paré (2011) Coarse root biomass allometric equations for *Abies balsamea*, *Picea mariana*, *Pinus banksiana*, and *Populus tremuloides* in the boreal forest of Ontario, Canada. *Biomass and Bioenergy* 135:4189-4196, <https://doi.org/10.1016/j.biombioe.2011.06.045>
- Brown S. (1997) Estimating Biomass and Biomass Change of Tropical Forests: A Primer. FAO Forestry Paper. Food and Agriculture Organization. Rome, Italy. 55 p.
- Brown S. (2002) Measuring carbon in forests: current status and future challenges. *Environmental Pollution* 116:363-372, [https://doi.org/10.1016/S0269-7491\(01\)00212-3](https://doi.org/10.1016/S0269-7491(01)00212-3)
- Cairns M. A., S. Brown, E. H. Helmer and G. A. Baumgardner (1997) Root biomass allocation in the world's upland forests. *Oecologia* 111:1-11, <https://doi.org/10.1007/s004420050201>
- Castellanos J. F., A. Velázquez M., J. Vargas H., C. Rodríguez F. y A. M. Fierros G. (1996) Producción de biomasa en un rodal de *Pinus patula*. *Agrociencia* 30:123-128.
- Chave J., M. Réjou-Méchain, A. Búrquez, E. Chidumayo, M. S. Colgan, W. B. C. Delitti, ... and G. Vieilledent (2014) Improved allometric models to estimate the aboveground biomass of tropical trees. *Global Change Biology* 20:3177-3190, <https://doi.org/10.1111/gcb.12629>
- Cole T. G. and J. J. Ewel (2006) Allometric equations for four valuable tropical tree species. *Forest Ecology and Management* 229:351-360, <https://doi.org/10.1016/j.foreco.2006.04.017>
- Díaz-Franco R., M. Acosta-Mireles, F. Carrillo-Anzures, E. Buendía-Rodríguez, E. Flores-Ayala y J. D. Etchevers-Barra (2007) Determinación de ecuaciones alométricas para estimar biomasa y carbono en *Pinus patula* Schl. et Cham. *Madera y Bosques* 13:25-34, <https://doi.org/10.21829/myb.2007.1311233>
- Djomo A. N., A. Ibrahimab, J. Saborowskic and G. Gravenhorst (2010) Allometric equations for biomass estimations in Cameroon and pan moist tropical equations including biomass data from Africa. *Forest Ecology and Management* 260:1873-1885, <https://doi.org/10.1016/j.foreco.2010.08.034>
- Domínguez-Cabrera G., O. A. Aguirre-Calderón, J. Jiménez-Pérez, R. Rodríguez-Laguna y J. A. Díaz-Balderas (2009) Biomasa aérea y factores de expansión de especies arbóreas en bosques del sur de Nuevo León. *Revista Chapingo, Serie Ciencias Forestales y del Ambiente* 15:59-64.
- Flores-Nieves P., M. Á. López-López, G. Ángeles-Pérez, M. L. de la Isla-Serrano y G. Calva-Vásquez (2011) Modelos para estimación y distribución de biomasa de *Abies religiosa* (Kunth) Schltld. et Cham. *Revista Mexicana de Ciencias Forestales* 2:9-20, <https://doi.org/10.29298/rmcf.v2i8.543>
- Fonseca W., F. Alice y J. M. Rey (2009) Modelos para estimar la biomasa de especies nativas en plantaciones y bosques secundarios en la zona Caribe de Costa Rica. *Bosque (Valdivia)* 30:36-47, <https://doi.org/10.4067/S0717-92002009000100006>
- Fonseca W., F. E. Alice and J. M. Rey-Benayas (2012) Carbon accumulation in aboveground and belowground biomass and soil of different age native forest plantations in the humid tropical lowlands of Costa Rica. *New Forests* 43:197-211, <https://doi.org/10.1007/s11056-011-9273-9>
- Gárate M. y J. A. Blanco (2013) Importancia de la caracterización de la biomasa de raíces en la simulación de ecosistemas forestales. *Ecosistemas* 22:66-73, <https://doi.org/10.7818/ECOS.2013.22-3.10>
- Grier C. C., K. A. Vogt, M. R. Keyes and R. L. Edmonds (1981) Biomass distribution and above- and below-ground production in young and mature *Abies amabilis* zone ecosystems of the Washington Cascades. *Canadian Journal of Forest Research* 11:155-167, <https://doi.org/10.1139/x81-021>
- Henry M., A. Bombelli, C. Trotta, A. Alessandrini, L. Birigazzi, G. Sola, ... and L. Saint-André (2013) GlobAllomeTree: international platform for tree allometric equations to support volume, biomass and carbon assessment. *iForest* 6:326-330, <https://doi.org/10.3832/ifor0901-006>
- IPCC, Intergovernmental Panel on Climate Change (2006) 2006 IPCC Guidelines for National Greenhouse Gas Inventories. H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.). IPCC National Greenhouse Gas Inventories Programme. Hayama, Japan. 20 p.
- Kenina L., A. Bardulis, R. Matisons, R. Kapostins and A. Jansons (2018) Belowground biomass models for young oligotrophic Scots pine stands in Latvia. *iForest* 11:206-211, <https://doi.org/10.3832/ifor2553-010>
- Ketterings Q. M., R. Coe, M. van Noordwijk and C. A. Palm (2001) Reducing uncertainty in the use of allometric biomass equations for predicting above-ground tree biomass in mixed secondary forests. *Forest Ecology and Management* 146:199-209, [https://doi.org/10.1016/S0378-1127\(00\)00460-6](https://doi.org/10.1016/S0378-1127(00)00460-6)
- Koala J., L. Sawadogo, P. Savadogo and E. Aynekulu (2017) Allometric equations for below-ground biomass of four key woody species in West African savanna-woodlands. *Silva Fennica* 51:1631, <https://doi.org/10.14214/sf.1631>
- Laclau P. (2003) Root biomass and carbon storage of ponderosa pine in a Northwest Patagonia plantation. *Forest Ecology and Management* 173:353-360, [https://doi.org/10.1016/S0378-1127\(02\)00012-9](https://doi.org/10.1016/S0378-1127(02)00012-9)
- Liepins J., A. Lazdins and K. Liepins (2018) Equations for estimating above- and belowground biomass of Norway spruce, Scots pine, birch spp. and European aspen in Latvia. *Scandinavian Journal of Forest Research* 33:58-70, <https://doi.org/10.1080/02827581.2017.1337923>
- Litton C. M., M. G. Ryan, D. B. Tinker and D. H. Knight (2003) Belowground and aboveground biomass in young postfire lodgepole pine forests of contrasting tree density. *Canadian Journal of Forest Research* 33:351-363, <https://doi.org/10.1139/x02-181>
- López L. M. A. (2005) Un procedimiento alternativo al tradicional para la medición de alturas con clinómetro. *Madera y Bosques* 11:69-77, <https://doi.org/10.21829/myb.2005.1121257>
- Magnani F., M. Mencuccini and J. Grace (2000) Age-related decline in

- stand productivity: the role of structural acclimation under hydraulic constraints. *Plant, Cell and Environment* 23:251-263, <https://doi.org/10.1046/j.1365-3040.2000.00537.x>
- Martínez D., S. Solís, F. J. Hernández y J. J. Graciano (2016) Ecuaciones para estimar biomasa aérea y subterránea en áreas reforestadas de *Pinus cooperi* Blanco en Durango, México. *Foresta Veracruzana* 18:25-32.
- Mugasha W. A., T. Eid, O. M. Bollandsås, R. E. Malimbwi, S. A. O. Chamshama, E. Zahabu and J. Z. Katani (2013) Allometric models for prediction of above- and belowground biomass of trees in the miombo woodlands of Tanzania. *Forest Ecology and Management* 310:87-101, <https://doi.org/10.1016/j.foreco.2013.08.003>
- Nam V. T., M. van Kuijk and N. P. R. Anten (2016) Allometric equations for aboveground and belowground biomass estimations in an evergreen forest in Vietnam. *PLoS ONE* 11:e0156827, <https://doi.org/10.1371/journal.pone.0156827>
- Návar J. (2015) Root stock biomass and productivity assessments of reforested pine stands in northern Mexico. *Forest Ecology and Management* 338:139-147, <https://doi.org/10.1016/j.foreco.2014.11.020>
- Ngoma J., E. Moors, B. Kruijt, J. H. Speer, R. Vinya, E. N. Chidumayo and R. Leemans (2018) Below and above-ground carbon distribution along a rainfall gradient. A case of the Zambezi teak forests, Zambia. *Acta Oecologica* 87:45-57, <https://doi.org/10.1016/j.actao.2018.02.003>
- Peichl M. and M. A. Arain (2007) Allometry and partitioning of above- and belowground tree biomass in an age-sequence of white pine forests. *Forest Ecology and Management* 253:68-80, <https://doi.org/10.1016/j.foreco.2007.07.003>
- Perry J. P. (1991) The Pines of México and Central America. Timber Press. Portland, Oregon, USA. 231 p.
- Picard N., E. Rutishauser, P. Ploton, A. Ngomanda and M. Henry (2015) Should tree biomass allometry be restricted to power models? *Forest Ecology and Management* 353:156-163, <https://doi.org/10.1016/j.foreco.2015.05.035>
- Picard N., L. Saint-André y M. Henry (2012) Manual de Construcción de Ecuaciones Alométricas para Estimar el Volumen y la Biomasa de los Árboles: del Trabajo de Campo a la Predicción. Organización de las Naciones Unidas para la Alimentación y la Agricultura y Centre de Coopération Internationale en Recherche Agronomique pour le Développement. Roma, Italia. Montpellier, Francia. 223 p.
- Rathore A. C., A. Kumar, J. M. S. Tomar, J. Jayaprakash, H. Mehta, R. Kaushal, ... and O. P. Chaturvedi (2018) Predictive models for biomass and carbon stock estimation in *Psidium guajava* on bouldery riverbed lands in North-Western Himalayas, India. *Agroforestry Systems* 92:171-182, <https://doi.org/10.1007/s10457-016-0023-z>
- Razakamanarivo R., A. Razakavololona, M. A. Razafindrakoto, G. Vieilledent and A. Albrecht (2012) Below-ground biomass production and allometric relationships of eucalyptus coppice plantation in the central highlands of Madagascar. *Biomass and Bioenergy* 45:1-10, <https://doi.org/10.1016/j.biombioe.2011.01.020>
- Ríos-Carrasco E., and J. J. Návar-Cháidez (2010) Root allometry of two subtropical plant communities of northeastern México. *Tropical and Subtropical Agroecosystems* 12:123-134.
- Sanquetta C. R., A. P. D. Corte and F. da Silva (2011) Biomass expansion factor and root-to-shoot ratio for *Pinus* in Brazil. *Carbon Balance and Management* 6:6, <https://doi.org/10.1186/1750-0680-6-6>
- SAS Institute (2011) SAS/STAT® 9.3 User's Guide. SAS Institute. Cary, North Carolina, USA. 8621 p.
- Schumacher F. X. and F. D. S. Hall (1933) Logarithmic expression of timber-tree volume. *Journal of Agricultural Research* 47:719-734.
- Tashi S., C. Keitel, B. Singh and M. Adams (2017) Allometric equations for biomass and carbon stocks of forests along an altitudinal gradient in the eastern Himalayas. *Forestry* 90:445-454, <https://doi.org/10.1093/forestry/cpx003>
- Ter-Mikaelian M. T. and M. D. Korzukhin (1997) Biomass equations for sixty-five North American tree species. *Forest Ecology and Management* 97:1-24, [https://doi.org/10.1016/S0378-1127\(97\)00019-4](https://doi.org/10.1016/S0378-1127(97)00019-4)
- Varik M., J. Aosaar, I. Ostonen, K. Lõhmus and V. Uri (2013) Carbon and nitrogen accumulation in belowground tree biomass in a chronosequence of silver birch stands. *Forest Ecology and Management* 302:62-70, <https://doi.org/10.1016/j.foreco.2013.03.033>
- Vela G. L. (1980) Contribución a la Ecología de *Pinus patula* Schl. et Cham. Publicación Especial No. 19. Instituto Nacional de Investigaciones Forestales. México, D. F. 109 p.
- Waring B. G. and J. S. Powers (2017) Overlooking what is underground: root:shoot ratios and coarse root allometric equations for tropical forests. *Forest Ecology and Management* 385:10-15, <https://doi.org/10.1016/j.foreco.2016.11.007>
- Watzlawick L. F., C. R. Sanquetta, A. A. Mello y J. E. Arce (2001) Ecuaciones de biomasa aérea en plantaciones de *Araucaria angustifolia* en el sur del estado de Paraná, Brasil. In: Simposio Internacional Medición y Monitoreo de la Captura de Carbono en Ecosistemas Forestales. J. Gayoso y R. Jandl (eds.). 18-20 octubre 2001. International Union of Forest Research Organizations. Valdivia, Chile. pp:1-10.
- Xiao C. W. and R. Ceulemans (2004) Allometric relationships for below- and aboveground biomass of young Scots pines. *Forest Ecology and Management* 203:177-186, <https://doi.org/10.1016/j.foreco.2004.07.062>
- Xiao C. W., J. C. Yuste, I. A. Janssens, P. Roskams, L. Nachtergale, A. Carrara, ... and R. Ceulemans (2003) Above- and belowground biomass and net primary production in a 73-year-old Scots pine forest. *Tree Physiology* 23:505-516, <https://doi.org/10.1093/treephys/23.8.505>
- Yuen J. Q., T. Fung and A. D. Ziegler (2016) Review of allometric equations for major land covers in SE Asia: uncertainty and implications for above- and below-ground carbon estimates. *Forest Ecology and Management* 360:323-340, <https://doi.org/10.1016/j.foreco.2015.09.016>
- Zapata-Cuatas M., C. A. Sierra and L. Alleman (2012) Probability distribution of allometric coefficients and Bayesian estimation of aboveground tree biomass. *Forest Ecology and Management* 277:173-179, <https://doi.org/10.1016/j.foreco.2012.04.030>